

Using technology and behavioral economics to promote development in the early years: The effect of an over-the-phone program

May 2024

Juanita Bloomfield, Ana Balsa, Alejandro Cid and Philip Oreopoulos¹

Summary

Early childhood in developing countries -where risk factors are more prevalent and resources more limited- demand effective and scalable models. We design and experimentally evaluate a telephonic assistance and messaging program for highly vulnerable families with children aged 0 to 3. The intervention focuses on supporting positive parenting practices at home, fostering language development, and offering one-to-one assistance for taking up the government benefits that families are entitled to. The program was implemented in Uruguay with 1360 families that qualified to receive support from the government agency Uruguay Crece Contigo. Treated families receive weekly calls and text and audio messages three times a week for 8 months. We find that the program increases the weekly frequency of parental involvement in stimulating activities and reduces parental stress. In addition, treated families achieve greater access to social benefits and programs including cash transfers and labor market programs. Effects on parental stress are larger for families that received cash transfers during the intervention, and effects on parental stimulation are larger by families experiencing higher economic shocks at baseline. (JEL: J13, I10, I20, H43)

Data availability

The datasets generated and/or analyzed during the current study are not publicly available due to the presence of protected information, but all de-identified datasets will be made available upon request from the corresponding author to replicate the results.

Code availability

All codes used to generate the results presented in this paper are available from the corresponding author on request.

Ethical Approval

For all the studies involved in the analysis, we received approval from the Universidad de Montevideo Institutional Review Board. The experiments complied with all relevant ethical regulations for work with human participants and involved informed consent of the parents, who were able to ask any questions they might have to a trained Research Assistant.

Previous registration

The full protocol is registered on the AEA RCT Registry, reference (AEARCTR-0008702).

¹ Juanita Bloomfield, j.bloomfield@um.edu.uy, Universidad de Montevideo; Ana Balsa, abalsa@um.edu.uy, Universidad de Montevideo; Alejandro Cid, acid@um.edu.uy, Universidad de Montevideo; Philip Oreopoulos, University of Toronto, philip.oreopoulos@utoronto.ca. We acknowledge financial support from the Inter-American Development Fund and Bernard Van Leer Foundation. We thank Paula Patrone and Sara Sant' Anna for excellent field supervision of the implementation of the program and research assistance. This paper was registered in the AEA Registry with RCT ID number AEARCTR-0008702. The project was approved by the Institutional Review Board at Universidad de Montevideo, resolution CE 2021/2.

I. Introduction

More than 200 million children do not reach their developmental potential due to poverty, poor nutrition, and poor care (Grantham-McGregor et al., 2007). Families of low socioeconomic status spend less time with their children, show them fewer expressions of affection (Guryan, Hurst and Kearney, 2008; Berlinski and Schady, 2015; Bradley et al., 2001) and are less likely to read and interact with them (Berlinski and Schady, 2015; Bradley et al., 2001). Developmental deficiencies in children from poor families are generated at an early age and have negative consequences for those children in the long term (Almond, Currie and Duque, 2018) as well as for the next generation (Doepke, Sorrenti and Zilibotti, 2019).

Existing literature demonstrates the efficacy of interventions targeting parenting competencies and early environments (Heckman, 2006; Walker et al., 2011; Gertler et al., 2014; Attanasio et al., 2022). For example, home visiting programs, geared towards enhancing parental attitudes and behaviors, have been shown to improve child development (Britto et al., 2017). However, such interventions remain scarce, often restricted to small cohorts, and in many cases, they are short-term and too expensive. The challenge is to design parenting interventions that can be scaled to large fractions of the population.

In this paper we show that an over-the-phone parenting intervention can be a cost-effective and scalable vehicle for nurturing parental competencies and strengthening early environments. The program combines weekly telephone encounters between the parent and a program facilitator, text and audio messaging, a chatbot, and an artificial intelligence tool that gives parents automatic feedback on their language interactions with their children. The intervention is aimed at vulnerable families with children between the ages of 0 and 3 that qualify to receive support from Uruguay Crece Contigo (UCC), the Uruguayan government agency in charge of early childhood protection.² It runs for 8 months, with a first module dealing with general parenting beliefs, attitudes and practices, and a second one focused on language stimulation.

The program includes different kinds of low-cost outreach efforts at once that could be easily scalable. Even though we cannot perfectly disentangle which component of the program drives the results, we believe that using a composite of components is a unique feature of this intervention as it increases the likelihood of effectively reaching families – i.e. if one type of

² Families that qualify to receive support from UCC present indicators of social vulnerability and possible health problems. On average, 31% of mothers that qualify for the program have completed middle school education, which is much lower than the population average (45% according among families with children aged 0-36 months according to the Uruguayan National Survey of Nutrition, Child development and Health, 2018).

intervention doesn't work or resonate with one family (but does with another) perhaps another type will.

We evaluate the program through a randomized controlled trial. We randomly divided 1360 families into a treatment group and a control group. The treatment group receives calls and text and audio messages for eight months, they can navigate freely through the chatbot and have access to the feedback tool. The control group is not exposed to teleassistance or messages, although it is exposed to a restricted version of the chatbot with information on local resources. The evaluation has two follow-up surveys: one four months after the launching of the intervention and another one after eight months. In this way, we see the impact of the intervention after applying the first thematic module and after applying the entire intervention. We compare the treatment group with the control group using an intention-to-treat analysis.

We evaluate families using a telephonic questionnaire and language metrics obtained from WhatsApp audios with parent-child conversations that families send after the intervention. From the questionnaire, we consider four families of outcomes: (i) access to government benefits and programs, (ii) parental involvement and knowledge, (iii) parental wellbeing and stress, and (iv) child's anthropometry. From WhatsApp audios, we extract measures for the number of vocalizations of the adult, number of vocalizations of the child, duration of vocalizations, number of conversational turns and pitch range in the adult's voice.

In the first follow-up, we find that the intervention increased families' access to government benefits (by 0.30 standard deviations), and labor programs promoted by the government. We find no effects on the frequency of reported parental involvement with the child.

In the second follow-up, after 8 months of intervention, we find that treated families increase the frequency of parental involvement on cognitive stimulation by 0.19 standard deviations (SDs) and increase parental knowledge on language development by 0.15 SDs. Parental wellbeing is higher for treated families (by 0.16 SDs), parental stress is lower (by 0.2 SDs), and the treatment group reports having greater access to government transfers. In addition, we find that average word rate per minute of the mother is higher among audios of treated families (by 0.37 SDs), as well as the mother's voice pitch (by 0.36 SDs).

We also explore some potential mechanisms behind the findings. Families with no initial exposition to one of the main government's cash transfers, the Uruguay Social Card, show larger responses to the intervention in terms of parental stress and wellbeing, and access to government's benefits. Moreover, the intervention seems to be more effective to increase parental involvement among families with more economic shocks at baseline. Finally, there are also heterogeneous effects by educational level, with effects for parental stimulation and stress being stronger for

families with higher education at baseline, which were those that resulted being more intensively treated.

This work is organized as follows. Section II describes the related literature. Section III has information about the program. Section IV describes the methodology of the impact evaluation. Section V describes the estimation method. Section VI shows the results and Section VII concludes.

II. Related Literature

Our paper contributes to a vast literature on the impacts of early childhood interventions on child development and to a growing literature on the benefits of using technology in these interventions.

1. Early childhood interventions in general –

Research shows that early environments can be substantially enhanced by programs that expand and increase the quality of pre-school care, and by interventions that foster parental competences. Evaluations of the long-run impacts of preschool programs, such as the Perry Preschool Program and Head Start, have shown positive effects on educational attainment and earnings (Currie, 2001; Heckman et al., 2010; Garces et al., 2002). In addition, numerous studies show the effectiveness of parental interventions. In developed countries, these programs have demonstrated positive impacts for children's development, ranging from 0.3 to 0.5 standard deviations, alongside enhancements in parental skills (Bakermans-Kranenburg, Van Ijzendoorn and Juffer, 2003; Blok et al., 2005; Al et al., 2014). Similarly, in developing countries, parental interventions have been effective to improve the parent-child relationship, the parents' knowledge of the child's development and the child's mental and motor development (Nores and Barnett, 2010; Baker-Henningham and López Bóo, 2010; Knerr, Gardner and Cluver, 2013).

Among parental interventions, home visits have been a particularly effective mechanism to intervene in low-income households. These interventions have been shown to improve parenting skills (Wilson, 2010), as well as children's cognitive abilities (Walker et al., 2015; Attanasio et al., 2014), socio-emotional abilities, and behavior (Pickering and Sanders, 2014). The main constraint of home visiting programs relates to scalability. These require an intensive use of qualified facilitators, as well as high levels of training and supervision to ensure adequate execution of the intervention protocols (Leer et al., 2016), which makes it difficult to scale them up.

2. Early-childhood interventions that use technology to support parents –

Early-childhood interventions that use technology to support parents are increasingly drawing attention due to their potential of expanding programs' reach at a low cost. In addition, technology has the potential to facilitate the adaptation, personalization and focalization of programs. Technology can be useful to screen families and adapt interventions to provide solutions to specific needs that families may have.

Recent text-messaging interventions for parents have found favorable impacts on parents' involvement and child literacy skills (York, Loeb and Doss (2019), Doss et al. (2017), Meuwissen et al. (2017), Mayer et al. (2018), Hurwitz et al. (2015)), alcohol consumption and maternal empowerment during pregnancy (Evans et al. (2012), and preschool attendance (Ajzenman et al. (2022)). In Bloomfield et al. (2022) and Balsa et al. (2021) we show that a positive parenting e-messaging program rooted on behavioral economics insights is effective at increasing the frequency of parental investment and the quality of the language interactions between the caregiver and the child.

Other related interventions use *automated calls*, meaning that caregivers receive 1–2-minute calls where they listen to a recording. In rural Guatemala, a 2-month intervention using automated calls for caregivers of children aged 6 to 33 months increased caregivers' interactions with their children (by 0.11 SDs) and improved children's vocabulary (by 0.22 SDs). The intervention had no impact on maternal anxiety or overall child development (Arteaga and Trias, 2023). In Jamaica, a program consisting of a parent manual and play materials, bi-weekly calls from community health workers, and weekly text messages for caregivers of children aged 5-to-24-months, increased interactions with children (by 0.34 SDs) and praise, but had no impact on play materials in the home or use of interactive language (Smith et al., 2023). In Jordan, phone calls from community health volunteers to caregivers of children aged 6-to-36 months reduced caregiver depression (by 0.11 SDs) but had no effect on their anxiety or self-efficacy or on children's behavior and development (Rafla et al., 2024). In India, an 6-month-long intervention consisting of automated calls, with duration of 1-2 minutes each, with parenting advice for children aged 6 to 30 months, counter to expectations, reduced caregivers' self-efficacy and increased their anxiety (Arteaga et al. 2024). The authors provide some reasons for the disappointing results: (i) the flexibility in the the pace to complete the program, (ii) the lack of differentiated activities by children's age, (iii) insufficient focus on child development, (iv) the absence of caregiver-child activities or games.

This paper contributes to the scarce but growing literature on the use of technology to scale-up evidence-based guidance on child-rearing practices in low- and middle-income countries. Our intervention combines 15-minute calls from a facilitator with a chatbot and e-messages to support parents and offer one-to-one assistance. Moreover, within early childhood interventions that use

technology, we are pioneers in the use of AI, by embedding to the program an AI tool that makes it possible to adapt and personalize the intervention to families' characteristics.

3. Importance of face-to-face assistance –

Previous literature has underscored the importance of face-to-face assistance within interventions. Gallego et al. (2023) make a strong case that personal support, guidance and motivation are critical for the effectiveness of interventions to meaningfully change life trajectories. While low-touch nudges may be cost-effective, these tools target short-term actions and fail to address deeper motivational factors that underpin routine behavior. In order to generate sustained positive behavior and to produce long-term benefits, more intensive interventions that leverage personal assistance are needed.

Our previous research also points in this direction. We found, in Bloomfield et al. (2022) and Balsa et al. (2021) and Balsa et al. (2023b), that an e-messaging program implemented after a parenting workshop is effective to improve the frequency and the quality of parenting practices. However, we found in Balsa et al. (2022) that when the program was implemented in the context of the COVID-19 pandemic, with no previous parenting workshop, results are no-longer significant. Even though several hypotheses could reconcile the findings from these studies, we believe that the fact that messages were totally unrelated to personal contact in our second experiment (Balsa et al. 2022) is an important explanation.

III. The Program: Teleassistance in Early Childhood and Positive Parenting text and audio messages

A. Background

Following the height of the SARS-Cov-2 pandemic, the demand for assistance and referrals to the main home visiting program administered by the Uruguayan government was larger than the program's capacity.³⁴ In response to this situation, the government designed a teleassistance program for families who were on the waiting list for the home visiting program but could not be visited at their homes.

Building on the teleassistance intervention that UCC was designing and exploiting the universal access to smartphones – and the extended use of WhatsApp – in Uruguay, even among poor

³ The *Programa de Acompañamiento Familiar* (PAF) or Family Support Program is administered by Uruguay Crece Contigo, the government agency in Uruguay in charge of early childhood protection, and part of the Ministry of Social Development.

⁴ See for example Balsa et al. 2023a for evidence of the effect of the COVID-19 pandemic on household's stressors and wellbeing in Uruguay

households, the program added a text and audio e-messaging component that had already been found to be effective in Uruguay (Bloomfield et al., 2022; Balsa et al., 2021). The program also included a chatbot and an AI feedback device. The design of the program involved developmental psychologists, social communicators, speech therapists and behavioral economists, among others, in coordination with the government team in charge of implementing the intervention.

B. Intervention contents

The program is separated into two thematic modules with a duration of four months each. The first module covers general positive parenting guidelines including: health, nutrition, protection, stimulation and caregiver's self-care. The program addresses four parental competencies defined by Gomez and Muñoz (2014): affective (parental sensitivity and good treatment), protective (safe environment, routines, and feeding), stimulating (playing and reading) and reflective (self-assessment of parenting and self-care). In Appendix A, Table A1 shows the parenting guidelines in module 1 and Table A2 shows some examples of the messages sent.

The first module also has a strong emphasis on providing families with access to government benefits and services. This is an important feature given that disadvantaged people do not always take-up the benefits for which they are eligible, and this may have detrimental consequences (Aizer, 2003; Currie, 2004). If the costs of gathering information (Bhargava and Manoli, 2015), completing an application process (Bettinger et al. 2012), or the stigma perceptions due to program participation (Currie, 2004) are too high, individuals will not apply, or there may be psychological frictions such as procrastination, inattention and aversion to the program complexity (Bhargava and Manoli, 2015) explaining the failure to take-up a government support program. In this sense, helping families remove program application barriers can have significant effects on program participation.

The second module covers language stimulation. The importance of language development during early childhood is well known. It predicts outcomes during youth, such as literacy skills and school success (Zimmerman et al., 2009; Rowe, 2012), as well as outcomes in adulthood, such as mental health and unemployment (Law et al., 2009). The calls and messages encourage parents to pay attention to the child's non-verbal cues, to describe them the environment and initiate conversations, to wait for the child to respond and engage in two-way conversations, to tell them stories and to sing and read to them (see Tables A3 and A4 in Appendix A).

C. Components of the intervention

a. Telephone calls

Teleoperators are expected to make weekly voice calls to each family, although the frequency is adjusted to the uniqueness of each family. The teleoperator is always the same for each family and tries reaching five times each family before considering it as an unreachable case.

Teleoperators articulate the different local resources to which families have access and provide families with advice and guidance on parenting, health and nutrition practices. The duration of the calls has an average of 13 minutes.

b. Messages

Each module includes 48 messages that are sent via WhatsApp in text and in audio. Messages are sent three times a week and are personalized according to the child's name, age, and gender, and caregiver's name and gender. In module 1 we personalize messages of the following topics: feeding, limits/safe sleep, play and reading. We adjust messages to two age groups: children younger than 12 months and older children. In module 2, messages are tailored to four age groups based on the child's stage of development: 6 to 12 months, 12 to 24 months, 24 to 36 months and more than 36 months.

Messages include information about the benefits of improving parenting practices, tips for carrying out practical activities at home, as well as motivational nudges.⁵ The contents of the messages seek to mitigate suboptimal decisions caused by present bias, parental cognitive fatigue, and the influence of negative identities.⁶ These biases threaten the caregiver-child attachment, prevent parents from investing optimally, and ultimately affect child development.

Based on behavioral economics theory, our messages: (i) remind parents about the benefits of engaging in positive parenting behavior in order to make these benefits more salient and tangible, especially for those inclined towards immediate rewards, and improve investment through activation and recall of prior knowledge; (ii) provide parents with suggestions of simple and concrete activities to help address inattention and decision fatigue by decomposing the complexity of parenting into simpler tasks; (iii) provide parents with self-care suggestions and techniques to help address inattention and decision fatigue by improving parental self-control and emotional

⁵ The average message length across all modules is 433 characters. For module 1, informative messages average 501 characters, activity proposals average 446 characters, and reflection messages average 422 characters. In module 2, informative messages have an average length of 399 characters, activity proposals average 313 characters, and reflection messages average 465 characters.

⁶ In the area of early childhood development, behavioral economics has identified the following key barriers to parental investment: a) present-bias – families with large discount rates are less likely to invest in activities that provide benefits only in the future; b) inattention and cognitive fatigue – vulnerability of poor families may reduce the idle cognitive capacity to make accurate decisions; c) Negative identities – the absence of self-esteem and self-confidence prevent parents' motivation to undertake child rearing.

regulation; (iv) encourage parents to continue trying, to rely on their own resources and on the support of others, to help parents overcome negative identities and strengthen parental self-efficacy.

The sequence of messages in module 1 is as follows:

Message 1 (Tuesday). Message with information on a parenting topic.

Message 2 (Wednesday). Proposal for an activity to do at home.

Message 3 (Saturday). Reflection on the topic and motivation to continue practicing at home.

The sequence of messages in module 2 is as follows:

Odd week:

Message 1 (Tuesday). Message with information on a parenting topic.

Message 2 (Wednesday). Proposal for an activity to do at home.

Message 3 (Saturday). Audio request.

Even week:

Message 1 (24 hours after sending audio). Feedback to the audio.

Message 2 (Saturday). Reflection and motivational message.

Message 3 (Sunday). Reflection.

Messages include text, images and emojis, and are also sent in audio format. Figure A1 in Appendix A shows how a message is displayed in WhatsApp.

c. Chatbot

In addition to calls and messages, a specific WhatsApp chatbot was designed for the intervention. This tool allows for a two-way flow of communication between the research team and families. Families get access to a drop-down menu with information on the messages they previously received, the local resources available to them, and the possibility to notify changes to their mobile phone or to register the phone number of a family member.

Control families have access to a restricted version of the chatbot that only includes information on local resources. Figure A2 in Appendix A shows how the drop-down menu is displayed in WhatsApp for treated families.

d. AI feedback tool

In module 2, we incorporate an artificial intelligence tool that allows to automatically decode WhatsApp audios and generate feedback messages for families. The literature on early language interventions in Developmental Psychology (Ferjan Ramírez et al. (2019, 2020); Leech et al. (2018, 2021); Leung et al. (2020); McGillion et al. (2017); Suskind et al. (2016)) show that providing parents with feedback, or coaching, regarding their linguistic inputs can enhance parent-child interactions.

The tool was created specifically for this project with the aim of decoding communication and language patterns between the caregiver and the child. It is an innovative and easily scalable device useful for monitoring the child's language development and encouraging language stimulating practices on the side of the caregiver.

The tool can store, decode and respond automatically to WhatsApp audios (see Peurey et al., 2023, for a description of the tool). Every two weeks the family is asked to send an audio-recording of a language interaction between the caregiver and the child. Families send these audios through the WhatsApp chatbot and the recordings are automatically stored and decoded by an AI software⁷ that identifies the number of vocalizations of the caregiver and the child, the duration of vocalizations, the number of conversational turns, and the caregiver's voice pitch frequency and range. Those metrics are compared to the latest audio sent by the family and are used to prepare a personalized feedback message, that congratulates the caregiver on the areas that have improved and suggests activities to work on for areas that have not improved. The feedback message is sent 24 hours after the family sends the audio.

IV. Impact evaluation: Methodology

A. Experimental design

a) Sample and randomization

Participants were recruited from a list of families waiting to get support from the UCC Family Support Program (PAF), of the Ministry of Social Development (MIDES). The total sample size is 1360 families (See Appendix B for details of the recruitment process).

Participating households were randomized in a stratified manner according to the child's age and mother's education into a treatment group and a control group.⁸ The treatment group receives

⁷ We use the DiviMe software. We previously used this software in Balsa et al. (2022) to obtain language interaction metrics from videos. In this project we embedded DiviMe within an algorithm that stores audios, runs the software and provides automatic feedback to users.

⁸ Initially, we intended to have two treatment groups—one with teleassistance only for 4 months and the other one with teleassistance during the full 8-month program. However, the Ministry of Health considered that 4 months were not

teleassistance and messages for a period of 8 months, access to a chatbot with parenting information and local resources, and access to a language feedback AI tool. The control group only receives access to a chatbot with local resources.

The result of the randomization is shown in Table 1. We implemented the intervention in three batches.⁹ Intervention batch 1 has 600 households, of which 399 are from the treatment group and 201 are from the control group. Intervention batch 2 has 426 households, of which 285 are from the treatment group. Intervention batch 3 has 334 households, of which 167 are from the treatment group.

Table 1: Sample size in each treatment group and intervention batch

Batch	Control	Treatment	Total
1	201	399	600
2	141	285	426
3	167	167	334
Total	509	851	1360

b) Implementation

Telephone calls were carried out by MIDES teleoperators with the official telephone number of the Ministry.

We created and validated a WhatsApp account to send messages. In addition, each of the texts went through a validation process by the WhatsApp service (these validations are mandatory because WhatsApp wants to avoid spam messages). We generated outgoing message campaigns through a web platform that allows personalized messages to each user with the name and gender of the caregiver.

We asked families to provide us with at least one mobile phone where they could be reached. We sent the messages to all the contact numbers we had for each family (mothers, fathers and other caregivers) to maximize the chances that the message would reach the family. For the 399 treated families in batch 1, we sent messages to 528 mobile phones. For the 285 treated families in batch 2, we sent messages to 373 mobile phone contacts. And for the 167 treated families in batch 3,

sufficient to support the families and we extended teleassistance to all families for the full 8 months. This resulted in a 2/3 to 1/3 distribution between treated and control groups.

⁹ When the start date of the program arrived, UCC did not have the operational capacity to assist all treatment families through teleassistance at the same time, so we decided to randomly define three batches of implementation. At the beginning of batch 1, the teleassistance team consisted of 18 teleoperators, 1 supervisor and 1 administrative assistant. As of December 31, 2023, the teleassistance team was composed of: 16 operators, 1 administrative, 2 supervisors and 1 coordinator.

we sent messages to 213 mobile phone contacts. Eighty-three percent of the mobiles that received messages belonged to a woman and 17% belonged to a man. In 70% of households only one female caregiver received messages, in 1% of households only one male caregiver received messages, and in 29% of households both male and female caregivers received messages.

The control group received a welcome message to the chatbot. This message was sent to 201 families with 243 mobile phones in batch 1, and to 141 families with 168 mobile phones in batch 2, and to 167 families with 232 phones in batch 3.

Each intervention module is made up of 50 messages in total: a welcome message, 48 messages sent three times a week for 16 weeks, and a farewell message. For batch 1, the implementation of module 1 began on May 2022, the implementation of batch 2 began on November 2022 and the implementation of batch 3 began on June 2023.¹⁰

c) Compliance

During the implementation, the Uruguay Crece Contigo team asked us to exclude some families from the program given that they had complex situations that had to be dealt with more intensive programs. From the 851 treated families, 8% were excluded from the calls and 9% were removed from the messages. 7% of families were excluded both from calls and messages. Families also had the possibility of opting out from the treatment. 12% of families opted out from the calls and 8% opted out from the messages. 7% of treated families opted out from calls and messages. In Appendix C, Tables C1 and C2 describe families that either opted out or were excluded from the program. Families that opt out have less children, are more likely to have only one child, and have less risk of depression. Families that were excluded from the program have younger children, are less likely to receive government assistance, less likely to live in the capital city of Montevideo, have older mothers. In addition, in families that are excluded, it is less likely that the baseline survey was answered by the mother of the child.

The number of answered calls is 13 on average (see Table 2). When splitting the sample by maternal education, the number of answered calls is 15.8 for families in which the mother

¹⁰ In batch 2, between week 6 and 7 of the program, there was a pause in the outgoing campaigns of messages. This was a specific request from the Uruguay Crece Contigo team because they understood that the dates between 12/24/22 and 1/10/23 (around Christmas and New Year celebrations) were sensitive for the families receiving the messages and it was best to stop the flow of messages during that period. Teleassistance calls were also stopped during this period. This is a measure that is also applied to home visits carried out by the Uruguay Crece Contigo team (“home visiting” is another program of UCC: no household in the teleassistance program nor in the control group participate in the “home visiting” program).

completed middle school and 12 for families with less maternal education. The number of answered calls among families with and without economic shocks is relatively more similar.

Table 2: Number of calls with response, overall, by mother’s education level, and by economic shocks

	Average	Mother did not complete middle education	Mother completed middle education	Household did not receive economic shock	Household receive economic shock
All batches	13	12	15.8	13	13.2
Batch 1	12	11.4	13.8	12.2	12.1
Batch 2	15	13.3	19.8	14.6	15.3
Batch 3	4	4.4	3.3	4.7	4.0

Approximately 76% of the dispatched messages were successfully received or viewed on the intended mobile phones. Additionally, within households where at least one member received the messages, this proportion increases to 82%.

Furthermore, surveys were distributed to gauge treated families’ satisfaction with the messages. Nearly all respondents (98%) expressed a positive perception of the messages, and 93% found the weekly frequency to be suitable. The response rate of the satisfaction survey was 40% in module 1 and 17% in module 2. All families were then asked about their satisfaction with the chatbot in the follow-up surveys, where we have higher response rates.

B. Evaluation Scales

In this paper, we analyze data from two follow-up surveys conducted at 4 and 8 months after initiating the intervention, along with baseline survey data. All surveys were administered by telephone, and participants received 300 Uruguayan pesos (USD 8¹¹) as incentive, in the form of a mobile phone recharge upon survey completion. The collected socioeconomic data encompasses information about: (i) access to government transfers and programs, (ii) parental investment and

¹¹ This is really a tiny amount of money and does not affect the budget of a family (as a comparison, the official minimum wage in Uruguay is 550 USD).

knowledge, (iii) parental wellbeing and stress and (iv) child's anthropometry. In addition, we asked participants to send us a WhatsApp audio with a conversation between the caregiver and the child. Families received the suggestion of recording the audio during mealtime or reading time. The minimum extension required was 2 minutes.

Table D1 (in Appendix D) provides a summary of the various scales employed in each follow-up and batch. Within access to government benefits, we have information on access to cash transfers such as Uruguay Social Card¹² and the *Bono Crianza* (Parenting Bonus). We also consider access to food baskets in the previous 30 days of the survey and an indicator for receiving free early childhood care vouchers for household with no access to early education in their neighborhoods. Finally, we consider access to home improvement benefits. We group all the latter outcomes in a single index that we construct using the first component of a principal component analysis.

We also have information on access to government programs such as labor market programs that have a 6-month duration.

Within parental investment we employ the StimQ index that assesses frequency and quality of parental involvement in stimulating activities. We construct an index of parental knowledge in childrearing as the total score in four parental knowledge general questions that are answered using a dichotomous measure (YES/NO). We also construct a parental knowledge in language stimulation index, using three specific questions in this topic that are answered in a dichotomous manner.

Regarding parental wellbeing, we use different instruments that we group in a single index using a principal component analysis. In particular, we group the risk from depression from the CES-D scale, the score in the WHO-5 instrument, and questions of the well-being scale of Mc Conkey (2000).

We also consider parental stress as measured by the Parenting Stress Index and child's anthropometric measures including weight-for-age and height-for-age. Finally, we include metrics from WhatsApp audios including number of vocalizations, vocalizations' durations, number of conversational turns, voice pitch and pitch range. These metrics are measured for the mother and for the child.

C. Attrition

¹² The Uruguay Social Card (TUS) is the means of payment for different monetary transfer benefits granted by the Ministry of Social Development to households and people in situations of extreme socio-economic vulnerability. The amount of the transfer varies (approx. from 30 to 200 USD) depending on the composition of the household.

Of the 1360 participant families (600 in batch 1, 426 in batch 2 and 334 in batch 3), 942 were surveyed (450 in batch 1, 272 in batch 2 and 220 in batch 3) in follow-up 1 and 753 (362 from batch 1, 236 from batch 2 and 155 from batch 3) at follow-up 2. 312 families (133 from batch 1, 104 from batch 2 and 75 from batch 3) sent the WhatsApp audio during the second follow-up. Table 3 shows the result of a regression of a dummy that takes a value of 1 if the family was surveyed on a treatment assignment dummy, fixed effects of batch and a constant. The first column corresponds to the first follow-up, the second column corresponds to the second follow-up and the third follow-up corresponds to the probability of sending the audio in the second follow-up. In the three cases, sample attrition was very similar among treatment and control families and the small difference that exists is not significant.

Table 3: Sample attrition

	Probability of completing telephonic survey		Probability of sending audio
	First follow-up	Second follow-up	Second follow-up
	(1)	(2)	(3)
Difference in the response rate of treated and controls	-0.004 (0.026)	0.025 (0.028)	-0.031 (0.024)
Batch 2	-0.111*** (0.029)	-0.049 (0.031)	0.023 (0.027)
Batch 3	-0.092*** (0.032)	-0.135*** (0.034)	-0.002 (0.029)
Control response rate	0.753*** (0.025)	0.587*** (0.027)	0.242*** (0.024)
N	1360	1360	1360

Notes: * p<.1, ** p<.05, *** p<.01.

D. Balance and Descriptive Statistics

Table 4 shows the mean, standard deviation, and number of observations for a series of sociodemographic variables of the child (Panel A), the caregiver (Panel B), and the household (Panel C) at baseline. In Tables E1, E2 and E3 in Appendix E we show, respectively, descriptive statistics for respondents of the first follow-up, second follow-up and those that sent the audio. The first three columns correspond to control observations, the second three columns correspond to treatment observations, and the last three columns show the difference between treated and controls. The children participating in the study are on average 26 months old, 51% of them are boys and 48% are the first child in the family. Less than 0.40% of the sample corresponds to cases of twins. 50% of children attend an educational center.

Almost all the respondents are women (99%). In 95% of the cases, the respondent is the mother and on average she is 28 years old. 25% of caregivers worked in the week prior to the survey. 31% of the sample completed basic secondary education (grades 7, 8 and 9). According to the score obtained on the CESD10 scale, the depression risk index ($CESD \geq 10$) is around 38%.¹³ The risk of depression increases if the household is single parent. We did not find a significant positive correlation between the risk of depression and the age of the mother or the number of children in the household. 28% of caregivers run out of internet on their phone every 15 days.

With respect to household variables, the number of members is around 5. The number of children in the home on average is 2. 96% of households receive assistance from the government. 55% receive the Uruguay Social Card (TUS) and 79% receive Family Allowances from the Equity Plan – monthly monetary transfer (from 25 to 300 USD) for vulnerable households with pregnant women, children, or individuals with disabilities. 68% of households are overcrowded and 84% of them faced a negative shock such as the death of a family member, large debts or legal problems in the 12 months prior to the survey. On average, the number of negative shocks received was 2. 33% of the households are from Montevideo (the capital city of Uruguay – half of the population of the country lives in Montevideo).

The last three columns of Table 3 show the difference between treated and controls on each covariate at baseline. We report the coefficient, standard error and sample size, of a regression of each covariate on the treatment assignment dummy, taking 1 if the household is treated and 0 if it is not. Of the 28 variables analyzed, first child is significant (it is higher for treated families). The difference is statistically significant at the 5% level. In a set of 28 variables, we should expect a statistical difference at the 5% between treated and controls in 1.4 of them. We found only one, which reflects a good balance.

¹³ This figure is consistent with the findings from Balsa et al. (2024), which shows that the prevalence of depressive symptoms among a sample of pregnant who attend public health care providers in Uruguay is 36%.

Table 4: Descriptive statistics at baseline

	Control			Treatment			Difference		
	N	Mean	s.d.	N	Mean	s.d.	N	Coefficient	s.e.
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
<i>Panel A: Child</i>									
Age (months)	509	25.105	13.012	851	27.127	12.086	1360	-0.102	(0.578)
Male	509	0.505	0.500	851	0.515	0.500	1360	0.009	(0.028)
First son	509	0.456	0.498	851	0.504	0.499	1360	0.058**	(0.028)
Twin	509	0.033	0.180	851	0.039	0.193	1360	0.000	(0.011)
Attends an early childhood center	509	0.517	0.500	851	0.485	0.499	1360	-0.018	(0.028)
Lives with mother	509	0.976	0.152	851	0.980	0.140	1360	0.003	(0.008)
Father or male caregiver in hhld	509	0.607	0.489	851	0.618	0.486	1360	0.006	(0.028)
<i>Panel B: Caregiver</i>									
Woman	509	0.994	0.077	851	0.988	0.108	1360	-0.007	(0.006)
Age (years)	509	27.179	7.526	851	28.381	35.150	1360	1.267	(1.598)
Works	509	0.254	0.435	851	0.247	0.430	1360	-0.009	(0.024)
Relationship=mother	509	0.953	0.212	851	0.951	0.217	1360	-0.001	(0.012)
Completed middle school	509	0.306	0.461	851	0.323	0.468	1360	0.020	(0.026)
Completed high school	509	0.071	0.257	851	0.051	0.219	1360	-0.021	(0.013)
Risk of depression	509	0.394	0.489	851	0.366	0.482	1360	-0.030	(0.027)
Frequency with no internet access = every week	509	0.327	0.410	851	0.326	0.405	1360	-0.000	(0.023)
Frequency with no internet access = once a month	509	0.151	0.318	851	0.149	0.314	1360	0.010	(0.018)
Frequency with no internet access = every 15 days	509	0.266	0.385	851	0.267	0.380	1360	-0.005	(0.022)

Frequency with no internet access = every week	509	0.256	0.380	851	0.258	0.376	1360	-0.005	(0.021)
<i>Panel C: Household</i>	509	0.327	0.410	851	0.326	0.405	1360	-0.000	(0.023)
Number of members	509	4.786	1.957	851	4.708	2.206	1360	0.041	(0.118)
Number of children	509	2.177	1.375	851	2.041	1.205	1360	-0.096	(0.072)
Overcrowding	509	0.689	0.463	851	0.678	0.467	1360	-0.007	(0.026)
Government assistance	509	0.964	0.185	851	0.953	0.212	1360	-0.007	(0.011)
Receives TUS	509	0.578	0.492	851	0.534	0.498	1360	-0.029	(0.028)
Receives AFAM-PE	509	0.794	0.403	851	0.783	0.412	1360	0.018	(0.023)
Experienced negative shock	509	0.848	0.356	851	0.842	0.363	1360	0.000	(0.020)
Number of negative shocks	509	2.177	1.570	851	2.083	1.489	1360	-0.052	(0.086)
Montevideo	509	0.325	0.468	851	0.338	0.473	1360	-0.036	(0.025)

Notes: * p<.1, ** p<.05, *** p<.01. We impute missing observations with the mean for each variable.

V. Estimation Method

We used ordinary least squares to estimate the effect of being assigned to the treatment (ITT_i = dummy that takes on the value of 1 if the family i was randomly assigned to treatment and 0 otherwise) on the outcome variable of interest, Y_i , and B_i which is a batch fixed effect.

$$Y_i = \beta_0 + \beta_1 ITT_i + B_i' \beta_2 + S_i' \beta_3 + X_i' \beta_4 + e_i \quad (1)$$

We also adjust Equation (1) subsequently for the following set of covariates: (i) stratum fixed effects (S_i includes 4 strata, given by the combinations of two age groups of the child and two groups of mother's education¹⁴), (ii) a set of covariates X_i that were unbalanced at baseline and (iii) a baseline measure of the outcome variable if available.¹⁵

In the estimates we use robust standard errors. For each family of outcomes, we adjust standard errors to account for multiple hypothesis testing using the Romano-Wolf correction (Clarke et al., 2020). Families of outcomes are marked with a heading in bold.

VI. Results

In this section we present the main results of the program. We first show results for the outcome variables obtained from the questionnaire including (i) access to government benefits, (ii) parental inputs, (iii) parental wellbeing and stress, and (iv) child's anthropometry. Following, we present results for language metrics obtained from WhatsApp audios. Finally, we assess heterogeneous effects by: maternal socioeconomic level, whether the household received economic shocks, whether the family received TUS at baseline and whether the child is the first of his or her family.

A. Main results

Table 5 shows estimates of intervention effects on several outcome variables.¹⁶ Estimates were obtained from a regression of each dependent variable on an indicator of treatment intention that takes on the value of 1 if the household is treated and 0 if it is a control. The first four columns correspond to the results of the first follow-up and columns (5)-(8) correspond to the results of the second follow-up. Columns (1) and (5) show the raw treatment-control differences and the

¹⁴ We defined a stratum according to whether the mother completed middle school or not. We define another stratum according to whether the child's age was greater or less than 18 months as of March 10, 2022.

¹⁵ We also estimated a variation of the specification of equation (1) with control variables that have been selected based on the method described in Belloni et al. (2014). The method consists of selecting the variables that best predict the treatment status with a LASSO regression, and then selecting the variables that best predict the outcome, with a LASSO regression as well. Our results are robust to this alternative specification.

¹⁶ This table contains main results of the intervention using summary indices. Results of the full set of outcomes assessed are reported in Appendix F.

coefficient's unadjusted standard error. Columns (2) and (6) show ITT effects after adjusting for stratum used in the randomization, and the first child dummy. Columns (3) and (7) add a control for the outcome at baseline, if available. In columns (4) and (8) we report the p-values corresponding to the outcome difference by ITT status reported in columns (1) and (5) when adjusting for multiple hypothesis testing. Families of outcomes are marked with a heading in bold and italics. We report the number of observations for each regression in straight parentheses. In Appendix G we report power calculations to detect an effect of 0.2 standard deviations in continuous outcomes and of 20% in dichotomous outcomes.

We find that in the first and second follow-up, treated families report having more access to government transfers than control families. The magnitude of the effect is of 0.30 standard deviations (standard deviations of each outcome are reported in Table D1 in the Appendix) and is significant at the 1% level. After the first four months of intervention, there is also an increase in access to labor programs sponsored by the government in 0.23 standard deviations. In the second follow-up, the program increased the frequency of parental involvement in stimulating activities as measured by the Stim-Q. The effect is of 0.19 standard deviations. We also find an effect on parental knowledge in language stimulation of 0.15 standard deviations. We find no impact on parental investment quality and on parental knowledge in general childrearing topics. Regarding parental wellbeing, we find a decrease in parental stress, which is significant in the second follow-up. The reduction is of 0.19 standard deviations. Finally, results on child's anthropometry are non-significant, but power to detect an effect on these outcomes is less than 50% (see Appendix G).

Table 5: Effects of the intervention on access to government transfers and programs, parental involvement, caregiver well-being and child's anthropometry. Coefficients in standard deviations.

	1st follow-up			2nd follow-up				
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
Panel A: Government transfers and programs								
Government transfers	0.296*** (0.068) [873]	0.306*** (0.066) [871]	0.300*** (0.059) [866]	0.010	0.301*** (0.076) [705]	0.329*** (0.074) [704]	0.337*** (0.069) [700]	0.010
Labor programs	0.230*** (0.071) [939]	0.232*** (0.072) [937]		0.010	0.074 (0.079) [747]	0.068 (0.079) [746]		0.594
Panel B: Parental investment and knowledge								
Parental Knowledge – Childrearing	0.097 (0.068) [932]	0.099 (0.067) [930]		0.010	0.029 (0.079) [733]	0.015 (0.078) [732]		0.713
Parental Knowledge – Language					0.150** (0.075) [730]	0.145* (0.075) [729]		0.099
StimQ - Frequency					0.187** (0.077) [740]	0.181** (0.076) [739]		0.059
StimQ - Quality					-0.082 (0.071) [740]	-0.069 (0.071) [739]		0.891

Panel C: Parental Wellbeing and Stress

Wellbeing Index	0.006 (0.094) [467]	0.013 (0.094) [466]		0.307	0.156** (0.079) [697]	0.156** (0.079) [696]		0.267
Parental Stress Index	-0.066 (0.091) [479]	-0.061 (0.092) [478]		0.059	-0.193** (0.078) [697]	-0.193** (0.078) [696]		0.218

Panel D: Child's anthropometry

Weight for age	0.186 (0.121) [329]	0.185 (0.123) [328]	-0.152 (0.155) [155]	0.673	0.116 (0.107) [389]	0.109 (0.108) [388]	0.029 (0.113) [269]	0.228
Height for age	-0.107 (0.128) [309]	-0.108 (0.130) [308]	-0.107 (0.166) [133]	0.663	-0.070 (0.106) [354]	-0.080 (0.106) [353]	-0.162 (0.127) [217]	0.762
Controls								
Strata and unbalanced vars	N	Y	Y		N	Y	Y	
Variable in baseline if available	N	N	Y		N	N	Y	

Notes: * p<.1, ** p<.05, *** p<.01. Rows depict different outcomes. Families of outcomes are identified by a heading in *italics*. Columns (1)-(4) and (5)-(8) report the ITT coefficient (outcome difference between ITT=1 and ITT=0) in the first and second follow-up respectively. Columns (2)-(3) and (5)-(7) report the ITT coefficient of specifications that include controls. We adjust for multiple hypothesis testing the p-values of the coefficients in columns (1) and (5) and report these in columns (4) and (8) respectively. Families of outcomes are marked with a heading in **bold** and *italics*. Number of observations are reported in straight parentheses.

Table 6 shows results obtained from the WhatsApp audios. We consider outcomes of the mother, of the child and of both of them together.¹⁷ The average word rate per minute of the mother increases and is significant at the 10% level. In addition, the pitch range of mothers also increases, and the effect has a magnitude of 0.762.

Table 6. Effects of the intervention on language outcomes. Coefficients in standard deviations.

	Mother's average word rate per minute	Mother's number of vocalizations per hour	Mother's average duration in milliseconds of vocalizations	Child's number of vocalizations per hour	Child's average duration in milliseconds of vocalizations	Mother's pitch range	Mother's pitch	Number of turn transitions between the child and the adult
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
Coefficient	0.369* (0.216)	0.114 (0.149)	-0.067 (0.173)	0.036 (0.171)	-0.066 (0.143)	-0.152 (0.176)	0.355* (0.185)	-0.064 (0.162)
N	123	123	123	123	122	123	123	123

Notes: * $p < .1$, ** $p < .05$, *** $p < .01$. Columns depict different outcomes. The table reports the ITT coefficient (outcome difference between ITT=1 and ITT=0) in the second follow-up for each outcome. We control for strata and for a dummy that indicates whether the child is the first of his or her family. We also control for the outcome at baseline. Number of observations are reported below each coefficient.

B. Heterogeneous impacts

To assess the mechanisms by which the intervention operates, we explore program heterogeneity across three dimensions: (i) mother's education level, (ii) presence of economic shocks in the household, (iii) recipient status of the main cash transfer (TUS) at baseline, and (iv) first child.¹⁸ Results are presented in Tables 7, 8, 9 and 10. The first three columns correspond to the first follow-up and the last three columns correspond to the second follow-up. Columns (1), (2), (4) and (5) shows the coefficients and standard errors from an OLS regression of the outcome in each row on the ITT main effect, the interaction between ITT status and the variable capturing the mechanism, the main effect of the mechanism, maternal education, a dummy for whether the household received an economic shock, and a dummy for whether the household received TUS at baseline and randomization strata. The table displays only the ITT main effect and the interaction of ITT with the mechanism. Table 7 shows that families in which the mother has

¹⁷ We account for the length of the audio, either by the definition of the outcome or by controlling by the duration of the audio.

¹⁸ Due to the limited number of observations in the language outcomes, we do not consider these in the heterogeneity analysis.

completed middle school derive some additional benefits from the intervention in terms of access to government transfers (see Column (2)), parental involvement in stimulating activities and parental stress index (see Column (5)). These results are in line with the fact that families in which the mother completed middle school have less issues of connectivity, change phone numbers less often and are therefore easier to contact and to provide the intervention.

Table 7: Heterogeneous impacts by mother's education level. Coefficients in standard deviations.

	1st follow-up			2nd follow-up		
	ITT	ITT x completed middle school	N	ITT	ITT x completed middle school	N
	(1)	(2)	(3)	(4)	(5)	(6)
Government transfers	0.209*** (0.079)	0.267* (0.146)	873	0.263*** (0.090)	0.143 (0.164)	705
Labor programs	0.210** (0.085)	0.049 (0.156)	939	0.125 (0.096)	-0.149 (0.169)	747
Parental Knowledge - Childrearing	0.120 (0.082)	-0.055 (0.136)	932	-0.056 (0.097)	0.247 (0.165)	733
Parental Knowledge - Language				0.161* (0.091)	-0.027 (0.158)	730
StimQ - Frequency				0.136 (0.097)	0.147 (0.152)	740
StimQ – Quality				-0.148 (0.092)	0.247* (0.138)	740
Wellbeing Index	0.007 (0.114)	0.009 (0.195)	467	0.187** (0.092)	-0.075 (0.173)	697
Parental Stress Index	-0.010 (0.115)	-0.173 (0.179)	479	-0.090 (0.099)	-0.317** (0.160)	697
Weight for age	0.232 (0.156)	-0.125 (0.257)	329	0.134 (0.137)	-0.126 (0.224)	389
Height for age	-0.111 (0.164)	0.019 (0.267)	309	-0.080 (0.134)	-0.012 (0.217)	354

Note: Columns (1), (2), (4) and (5) denote a specific model in which the ITT is interacted with a dummy that takes value of 1 if the mother completed middle school. The first three columns correspond to the first follow-up and the second three correspond to the second follow-up. Each row denotes a different outcome. For each outcome, we show the coefficient and standard error for the ITT main effect, and the coefficient and standard error for the interaction between ITT and the behavioral barrier analyzed. Standard errors are reported in parentheses. All regressions adjust for randomization strata, maternal education, a dummy for whether the household received an economic shock and a dummy for whether the household received TUS at baseline. N corresponds to number of observations. *p<.1, **p<.05, ***p<.01.

Table 8 shows that families that experienced an economic shock at baseline, are more likely to increase the frequency of parental involvement in stimulating activities and the wellbeing index in the second follow-up (see Column (5)). In Bloomfield et al. (2023) we show that families with more negative shocks in the past have stronger behavioral barriers and engage less frequency with their children. Because the intervention, and in particular the messaging component, was designed to address these barriers, the results from Table 8 suggest that the intervention triggered the right channels. The heterogeneous effects for these families seem to appear after 8 months of intervention, but not earlier. In fact, there is a negative effect on the interaction coefficient for parental knowledge when considering the first follow-up.

Table 8: Heterogeneous impacts by economic shocks. Coefficients in standard deviations.

	1st follow-up			2nd follow-up		
	ITT	ITT x shocks	N	ITT	ITT x shocks	N
	(1)	(2)	(3)	(4)	(5)	(6)
Government transfers	0.494*	-0.196	205	0.354**	-0.063	705
	(0.271)	(0.308)		(0.148)	(0.172)	
Labor programs	-0.207	0.537	219	0.195	-0.161	747
	(0.321)	(0.368)		(0.133)	(0.163)	
Parental Knowledge - Childrearing	0.309	-0.576*	218	-0.088	0.155	733
	(0.250)	(0.293)		(0.158)	(0.180)	
Parental Knowledge - Language				0.242*	-0.123	730
				(0.133)	(0.160)	
StimQ - Frequency				-0.120	0.419**	740
				(0.154)	(0.176)	
StimQ - Quality				0.010	-0.109	740
				(0.134)	(0.159)	
Wellbeing Index	-0.065	-0.090	208	-0.051	0.293*	697
	(0.282)	(0.319)		(0.141)	(0.169)	
Parental Stress Index	0.032	0.183	215	-0.176	-0.025	697
	(0.293)	(0.331)		(0.137)	(0.166)	
Weight for age	0.415	0.030	167	-0.074	0.227	389
	(0.352)	(0.413)		(0.189)	(0.231)	
Height for age	0.296	-0.505	154	-0.258	0.244	354
	(0.404)	(0.455)		(0.190)	(0.232)	

Note: Columns (1), (2), (4) and (5) denote a specific model in which the ITT is interacted with a dummy that takes value of 1 if the family received an economic shock at baseline. The first three columns correspond to the first follow-up and the second three correspond to the second follow-up. Each row denotes a different outcome. For each outcome, we show the coefficient and standard error for the ITT main effect, and the coefficient and standard error for the interaction between ITT and the behavioral barrier analyzed. Standard errors are reported in parentheses. All regressions adjust for randomization strata, maternal education, a dummy for whether the household received an economic shock and a dummy for whether the household received TUS at baseline. N corresponds to number of observations. *p<.1, **p<.05, ***p<.01.

In Table 9, we interact the ITT dummy with a dummy that takes value of 1 if the family did not have TUS at baseline. As expected, we find that families that did not receive Uruguay Social Card before the intervention are more likely to receive government transfers during the intervention. The effect is persistent both in the first and the second follow-up. In addition, we find that having no transfers at baseline, and therefore receiving them during the intervention, increases the effect of the program on the wellbeing index and on parental stress. We find no heterogeneous effects on parental involvement. Taken together, this evidence suggests that the monetary component of the intervention seems to be the mechanism behind the effect on parental stress.

Table 9: Heterogeneous effects by no-TUS at baseline. Coefficients in standard deviations.

	1st follow-up			2nd follow-up		
	ITT	ITT x no TUS	N	ITT	ITT x no TUS	N
	(1)	(2)	(3)	(4)	(5)	(6)
Government transfers	0.086 (0.068)	0.473*** (0.121)	868	0.122 (0.083)	0.453*** (0.140)	701
Labor programs	0.156 (0.098)	0.155 (0.143)	934	0.173 (0.109)	-0.209 (0.158)	743
Parental Knowledge - Childrearing	0.119 (0.091)	-0.047 (0.132)	927	0.002 (0.105)	0.050 (0.156)	729
Parental Knowledge - Language				0.087 (0.100)	0.124 (0.151)	726
StimQ - Frequency				0.075 (0.102)	0.245 (0.151)	736
StimQ - Quality				-0.090 (0.093)	0.055 (0.145)	736
Wellbeing Index	-0.149 (0.123)	0.333* (0.188)	464	0.085 (0.105)	0.164 (0.158)	693
Parental Stress Index	0.090 (0.127)	-0.333* (0.181)	476	-0.033 (0.106)	-0.368** (0.159)	693
Weight for age	0.286 (0.175)	-0.201 (0.252)	328	0.107 (0.139)	-0.035 (0.221)	387
Height for age	0.076 (0.168)	-0.358 (0.263)	308	0.062 (0.141)	-0.321 (0.214)	353

Note: Columns (1), (2), (4) and (5) denote a specific model in which the ITT is interacted with a dummy that takes value of 1 if the family did not receive TUS at baseline. The first three columns correspond to the first follow-up and the second three correspond to the second follow-up. Each row denotes a different outcome. For each outcome, we show the coefficient and standard error for the ITT main effect, and the coefficient and standard error for the interaction between ITT and the behavioral barrier analyzed. Standard errors are reported in parentheses. All regressions adjust for randomization strata, maternal education, a dummy for whether the household received an economic shock and a dummy for whether the household received TUS at baseline. N corresponds to number of observations. *p<.1, **p<.05, ***p<.01.

Table 10 shows heterogeneous effects by a dummy that indicates whether the child is the first of his or her family. Firstborns show larger effects on government transfers, but smaller effects on parental knowledge in childrearing in the first follow-up. In the second follow-up there are differential effects on the wellbeing index, suggesting that families that received the intervention for their first child, have a higher wellbeing than those that have older children also.

Table 10: Heterogeneous effects by first child. Coefficients in standard deviations.

	1st follow-up			2nd follow-up		
	ITT	ITT x first child	N	ITT	ITT x first child	N
	(1)	(2)	(3)	(4)	(5)	(6)
Government transfers	0.192** (0.085)	0.244* (0.134)	871	0.227** (0.097)	0.211 (0.149)	704
Labor programs	0.289*** (0.096)	-0.147 (0.143)	937	0.182* (0.104)	-0.247 (0.158)	746
Parental Knowledge - Childrearing	0.247*** (0.088)	-0.317** (0.134)	930	0.075 (0.110)	-0.123 (0.155)	732
Parental Knowledge - Language				0.059 (0.096)	0.206 (0.151)	729
StimQ - Frequency				0.279*** (0.107)	-0.211 (0.151)	739
StimQ - Quality				-0.069 (0.101)	0.009 (0.144)	739
Wellbeing Index	-0.033 (0.129)	0.086 (0.189)	466	-0.056 (0.103)	0.478*** (0.156)	696
Parental Stress Index	0.052 (0.126)	-0.252 (0.182)	478	-0.117 (0.106)	-0.174 (0.157)	696
Weight for age	0.091 (0.171)	0.183 (0.244)	328	0.153 (0.150)	-0.138 (0.218)	388
Height for age	-0.057 (0.190)	-0.112 (0.258)	308	0.034 (0.145)	-0.269 (0.216)	353

Note: Columns (1), (2), (4) and (5) denote a specific model in which the ITT is interacted with a dummy that takes value of 1 if the child is the first from his or her family. The first three columns correspond to the first follow-up and the second three correspond to the second follow-up. Each row denotes a different outcome. For each outcome, we show the coefficient and standard error for the ITT main effect, and the coefficient and standard error for the interaction between ITT and the behavioral barrier analyzed.

Standard errors are reported in parentheses. All regressions adjust for randomization strata, maternal education, a dummy for whether the household received an economic shock and a dummy for whether the household received TUS at baseline. N corresponds to number of observations. *p<.1, **p<.05, ***p<.01.

VII. Conclusions

In this work we design, implement, and experimentally evaluate a teleassistance and telephone messaging program for families with children from 0 to 3 years old. The program was implemented in Uruguay with 1360 families that qualified to receive support from the government agency Uruguay Crece Contigo. The intervention focuses on supporting positive parenting practices at home, fostering language development and offers one-to-one assistance for taking up the government benefits that families are entitled to.

Overall, the program was well-received by treated families: 90% of families report that the chatbot is useful to gather information on parenting practices and language stimulation, and 75% of families report the chatbot is useful to gather information on local resources. In addition, teleoperators value the tool. From qualitative surveys, teleoperators report that messages allow to “break the ice” when calling families and anchor the sequence of calls around topics with which the family has already been in contact during the week. Also in some cases, facilitators prefer to start the conversation “chatting” instead of talking, which allows them to gradually gain the trust of the caregivers. An additional advantage is that with a telephone intervention it is possible to reach rural places, where there are no territory teams that can make home visits.

After 4 months of intervention, we find that treated families have more access to social benefits. The effects are of a magnitude of 0.30 standard deviations. Treated families have also more access to government sponsored labor programs (in 0.23 standard deviations).

In the second follow-up, after 8 months of intervention, parental involvement in stimulating activities, measured by questions of the StimQ, is 0.19 standard deviations higher for the treated group and parental knowledge in language stimulation also increases. In addition, parental stress is lower and wellbeing is higher for treated families (by 0.19 and 0.16 standard deviations respectively) and the treatment group continues to have more access to social benefits.

It is possible that one of the main channels through which the program affects parental involvement is by decreasing parental stress. Targeting parental stress seems to be key such vulnerable households where 68% of them are overcrowded and 84% faced negative shocks such as the death of family member, large debts or legal problems in the prior 12 months of the survey (see Table 4).

The lower parental stress achieved by the intervention may be the result of the helping hands of teleoperators that encourage the take up of social protection (we find greater access to monetary transfers and public programs). Indeed, we find heterogeneous effects depending on whether the household received or not received the Uruguay Social Card transfer at baseline. Parental stress

is lower for those that did not have this transfer before the intervention. The usefulness of the chatbot for parenting issues also seems to be important, as families that were more intensively treated with the intervention (those with higher educational level), and those with more behavioral barriers at baseline have higher impacts on the frequency of parental stimulation.

Since the results come from self-reported surveys, it is not possible to rule out the presence of an experimenter demand effect. However, the possibility of experimenter demand effect is undermined by the fact that questionnaires are not applied by the teleoperator -that may affect the answers of the families in the follow-up- but by external social workers that do not know the treatment group of the families. Finally, it should also be noted that families were instructed during the surveys that their answers would remain anonymous and would only be analyzed in an aggregate manner. This information may have likely mitigated the incentives for parents to offer the desirable answers.

Our results are in line with those from Bloomfield et al. (2023) and Balsa et al. (2023) in the sense that we find that the intervention is useful to increase parental involvement and improve language interactions between the parent and the child. In this paper, we additionally show that parental stress and wellbeing can be influenced by including one-to-one telephonic assistance to obtain cash transfers that families are entitled to receive.

The next version of this work will complement the results from parental questionnaires with administrative registries from the Ministry of Social Development. We are also planning a third follow-up in families' homes to assess child development.

Appendix A: Intervention contents

Table A1: Parenting topics included in intervention module 1.

Week	Topic	Parental competence
1	Parental sensitivity	Affective
2	Parental sensitivity	Affective
3	Good treatment	Affective
4	Safety at home	Protective
5	Routines	Protective
6	Routines	Protective
7	Feeding	Protective
8	Feeding	Protective
9	Limits / Safe sleep	Protective
10	Play	Stimulating
11	Play	Stimulating
12	Reading	Stimulating
13	Reading	Stimulating
14	Self-care	Reflective
15	Self-care	Reflective
16	Parental involvement	All

Table A2: Examples of messages sent during module 1, by topic and parental competence addressed

Topic	Parental Competence	Type of message	Message
Parental Sensitivity	Affective	Informative	When (child's name) is fussy, do you notice that you understand him or her better when you put yourself in his or her place? Their face will give you clues about what they need, so trust your intuition! Try different responses like giving him/her a hug or sitting on your lap. And even if you're wrong sometimes, know that the most important thing for your child to feel safe is to be attentive and always try to respond sensitively. 🧡
		Activity proposal	We all have difficult days, when we are more tired and less patient. On those days, if (child's name) cries or is fussy, try to manage your discomfort and mood so you don't get angry with (her). You can talk to her and express what is happening to you. Try saying, "Mommy is tired today. Or ask her "why are you crying, how can I help you?" Talking to her in a calm and loving tone will help you tune in to (child's name) even on those days when it's hard. 😊
		Reflection	These past few weeks we've been seeing that when (child's name) cries or is fussy, he/she may want to tell you a lot of things. It's normal to be worried, upset or upset, especially when she seems inconsolable. She's just trying to tell you what's wrong. Keep exercising your watchful eye and when you are overwhelmed, be sure to turn to other adults around you.
Good treatment Good treatment	Affective	Informative	Children who feel loved and contained develop and learn better. It is not necessary to set aside long hours, (child's name) will feel your love and containment as long as you are emotionally willing and give yourself in the daily moments of shared enjoyment. That is the sure foundation that will make your (daughter) feel happy and loved.

			Also, there's something you need to know: there's never too much cuddling! ❤️ There's a false belief that cuddling is a habit, but it's not. Don't hesitate to cuddle (child's name) when you see (child's name) looking for it.
		Activity proposal	Children need to be treated with care and kindness. Always avoid sudden movements when dealing with (child's name). Talk softly and never shake (child's name) to stop crying. When (child's name) cries, you can soothe him/her by holding him/her or putting him/her very close to you. You'll find it works and it's a way to show your unconditional love - we all like to be treated well! 😊
		Reflection	Your love is extremely important to your child's development. We know you want the best for (child's name) and (she) feels it. Don't worry if some days you feel distant or disconnected from (her). When we feel stressed this can be normal. At those times, find a way to relax and lean on someone else to take over (child's name)'s care. When you are calmer, you will feel connected again. Trust the value of hugs and kisses. Never put (child) to bed without first giving him/her a hug. ❤️
Routines	Protective	Informative	Routines give children security and confidence and also organize family life. To generate them, we must organize daily activities (such as feeding, sleeping and bathing) and repeat them. The more we repeat them, the stronger the routine becomes established. Toddlers don't understand times or schedules. But (child's name) will gradually understand "what comes next" if you repeat his daily routines of feeding, bathing and sleeping in the order that works best in your home. What are (child's name)'s routines? 📅 ⌚ "
		Activity proposal	Games add a lot on an emotional level and you can include them in the day's activities to help (child's name) anticipate routines. Sometimes it helps to repeat the same song at bath time (remember "Al agua pato"?) or look for the same toy before taking (child's name) to the bathroom. 💧 That way, we are sure that (child's name) will enjoy the bath. Do you have a bedtime song? 😊 Here we leave you some ideas: (link to a page with song playlist)
		Reflection	Do you notice changes in (child's name) when you struggle to follow the same routine every day? Routines provide containment, security and lower anxiety - and that helps both (her) and you enjoy it so much more! 😊 How about incorporating a bedtime story into your sleep routine?
Play	Stimulating	Informative	As she gets older, (child's name) is likely to become more fidgety and find it hard to stay with the same toy for long periods of time. She'll play with you sometimes, other times she'll choose to play alone, and she'll probably have trouble sharing her things. And she will probably find it hard to share her things, which is normal for her age!

			To teach him to share, try telling him that if a friend borrows a toy, he won't take it away. Sharing is learned and you have to help him/her do it. 
		Activity proposal	We know (child's name) is very active at this time. It's age. It's normal for her to seek new challenges with her body like jumping, climbing or running. She will always feel safer if you are close by. This week we invite you to explore objects and put toys together with him/her. Here's an idea: Find a string you have at home and two or three rolls of toilet paper (without paper). It's a worm! You can paint it or decorate it together. 
		Reflection	At this stage when (child's name) is so wiggly, we want to remind you that screens (TV, cell phone, computer) are not a recommended option for children under 4 years old. They provide a lot of information that (child's name) cannot process yet and, if you think about it, being in front of the screen takes time away from things that do add up, such as playing and communicating with other people. But if you need to resort to them at certain times, be sure to control the quality of what she sees and interact with her while she is in front of the screen. The benefits of limiting or even eliminating screen time in these early moments will last a lifetime. 
Reading	Stimulating	Informative	Reading and stories, as well as play, also help (child's name) learn and develop. Reading gives her vocabulary, helps her imagination and intelligence unfold. Stories envelop children in magical worlds full of emotions. Tell stories, change the tone of voice, immerse (her) in that world. (Child's name) will play an increasingly active role as you read to her. He/she will ask questions, interrupt your stories. Let him (her) interact with the books, he (she) may want to see only the illustrations or start at the end. Whatever exchange you choose to make with the book, it will allow him/her to get closer to the reading.
		Activity proposal	Reading brings a lot to people. The best thing to do is to include it in the daily routine and, in addition, to enjoy it. We invite you to think about what quiet time would be a good time to read to (child's name). We believe that the time before bedtime is, in general, a good time because it helps your child to calm down before going to sleep. Sometimes it's not necessary to have a book. You can choose a magazine, show him/her pictures and create simple stories, help him/her to imagine! Try asking him questions. For example: "What do you think is going to happen next? How many animals are there? What color are their eyes? Let him/her participate in the stories, changing the names of characters for family names. If he/she gets involved with reading and stories, he/she will develop more his/her language, imagination and intelligence.  
		Reflection	Reading time can become a space for connection and daily enjoyment shared by both of us. Take advantage of reading to him/her every day. Transform that time into something special for both of you. Because you are also giving him/her the pleasure of reading and that will last a lifetime.

			Enjoy every book together, every shared story, celebrate the stories! 📖👶
Self-care	Reflective	Informative	There are emotions such as anger, sadness or fear that, if they accumulate, affect our health. In addition, they influence our relationships. That is why it is very important to realize how you feel, to find a good way to manage your emotions when you are not well. Breathing, going for a walk or simply leaving the place for a few minutes when you are overwhelmed, are things that serve to regulate emotions. And don't hesitate to ask for help from someone you trust if you need it. Parenting is demanding, but it's also beautiful. 🌈
		Activity proposal	If you are stressed, overwhelmed or feel back pain, then we invite you to follow some of these tips. Take a deep breath. Take a break from whatever you are doing. Drink a glass of water. Take a minute to drink mate. All these practices generally work to lower your blood pressure. You have to find out what works best for you. Something that also helps to loosen tensions are gentle movements with the head. First moving it up and down as if saying "yes". Then to the sides as if saying "no". And after that, try moving your shoulders up and down as if saying "I don't know". If you repeat this several times, your neck will be much more relaxed. And of course, you can always go to your health center for a consultation if you feel persistently overwhelmed.
		Reflection	Try to express your feelings and emotions by saying "I'm nervous", "I feel exhausted today". Putting emotions into words generates relief. Remember that breaks, breathing, walking, or neck and head exercises help improve your mood and stress level when you feel overwhelmed or overwhelmed. Have you tried any of these suggestions and how did they go? The more relaxed you feel, the more you will enjoy your time with (child's name). 🧘💙

Table A3: Language stimulation topics included in intervention module 2.

Week	Topic	Parental competence
1-2	I understand more than I can express	Stimulating
3-4	Between gestures and words	Stimulating
5-6	Conversational turns	Stimulating
7-8	Feeding time	Stimulating
9-10	Verbal productions	Stimulating
11-12	Sentences	Stimulating
13-14	Music time	Stimulating
15-16	Reading	Stimulating

Table A4: Examples of messages sent during module 2, by topic and age of the child

Topic	Child's age	Type of message	Message
I understand more than I can express	12-24 months	Informative	Your child understands much more than he/she can express. At this stage, (CHILD'S NAME) clearly understands simple words and commands (e.g., "(CHILD'S NAME), Can you bring me your shoes, please?"). It is important for the child's language development that you ask him/her to reach for objects or point to others whenever possible. Also try to give him simple commands and ask him simple questions about things happening around him. 😊
		Activity proposal	Today we suggest that at some point in your play with (CHILD'S NAME), you pay attention to what he/she understands. For example: you can ask her to bring you the toy box; to take the ball out of the box; to give you a book; or to show you where the doll is. 🍀🧸
		Reflection	Have you been able to notice the words and commands that (CHILD'S NAME) understands? Language is not only what children express, it is also what they understand. By challenging him with increasingly complex questions and requests, you encourage (CHILD'S NAME) to continue to develop his understanding and language acquisition. Incorporate these challenges into everyday life as if they were games - both you and (CHILD'S NAME) will enjoy it! 😊
Between gestures and words	24-36 months	Informative	At this stage of development, (CHILD'S NAME) communicates a lot through his body language, with the gestures of his face, the movement of his shoulders, his hands, arms and postures in general. At this time, it is important that you accompany their body communication by putting into words what you think they are communicating. In this way, you are expanding their language. 💬❤️
		Activity proposal	Today, while you are feeding (CHILD'S NAME), pay attention to all the ways she uses to communicate. Try to put into words what you think he wants to communicate and wait for his response before you speak again. For example, (CHILD'S NAME) points to her glass on the table. You can say, "that's your glass, are you thirsty?"; (CHILD'S NAME) says "yes" and you say "here you go."
		Reflection	Were you able to identify what (CHILD'S NAME) was communicating to you in its own way during the week? This process doesn't happen overnight, it builds little by little. 🤝 By expressing out loud what you think (CHILD'S NAME) wants to say, you make your child feel recognized and understood, and you help him/her to incorporate the language better. 😊 Keep looking for these shared moments of communication and discovery with (CHILD'S NAME)!

Conversational turns	36+ months	Informative	In communication with (CHILD’S NAME) it is important to give space: that you listen and respond to what (CHILD’S NAME) wants to tell you and that you give space for him/her to respond to what you tell him/her. Communication is richer when it happens in turns, like in ping-pong: one serves, the other receives and returns. 🤝 When you manage to keep more turns of conversation with (CHILD’S NAME), you are stimulating their brain development: their intelligence, language acquisition, social and emotional skills. 💡
		Activity proposal	We propose that today, at some playtime with (CHILD’S NAME), you talk to him/her about what they are playing. 🍀🧸 Try to put yourself at his/her level, sit on the floor with him/her. While you are talking to him/her, try to leave space for him/her to express him/herself in some way and then continue talking to him/her. Remember that talking to (CHILD’S NAME) in turns is the best way to promote his/her language. practice it during this week! 😊
		Reflection	How did you do this week with the conversational turns between you and (CHILD’S NAME)? A good conversation has spaces for listening and moments of response. In order for children to express themselves, it is important to give them space to do so 😊. Remember this whenever you're talking to (CHILD’S NAME). This way you will be respecting their intention to communicate and promoting their language development.
Feeding time	6-12 months	Informative	Feeding time is the best space for communication. It allows physical, visual and auditory contact between (CHILD’S NAME) and you. Take advantage of these moments to interact with (CHILD’S NAME) and share words related to the foods he/she likes, dislikes, colors, sizes, textures, sweet-salty, etc. This way you are filling their world with words. 🍷👄 We share a link [LINK]
		Activity proposal	Today, when you are feeding (CHILD’S NAME), tell her who made the food, if it's warm or hot, soft, etc. 😊
		Reflection	How did it go this week conversing with (CHILD’S NAME)? Conversing with your child at feeding times creates rituals for life. 🍷 In those moments of interaction, avoid distractions such as TV, tablet or cell phones that hinder the face-to-face encounter with your child. Conversing with (CHILD’S NAME) makes him/her feel understood and loved. keep looking for these moments!

Verbal productions	24-36 months	Informative	(CHILD’S NAME) is acquiring new words every day. While you may not yet understand everything your child says, his or her everyday vocabulary is growing significantly. It is important that you put in your words and add new words to what (CHILD’S NAME) points, looks and/or says. In this way you enrich their vocabulary. 🗣️
		Activity proposal	Today we suggest that, at feeding time, you pay attention to the words that (CHILD’S NAME) uses to communicate with you and try, based on them, to add new words to his/her language. For example: (CHILD’S NAME) points and says "ota", you say: "Do you want the ball?" ... you wait for an answer from (CHILD’S NAME) and then you add more words to the one he/she has: "Ok, let's go get the ball that is on top of the chair!" ⚽👤
		Reflection	How did you do this week putting into words what (CHILD’S NAME) communicates to you and adding new words to her vocabulary? To accompany the language acquisition process, it is important to put into words what interests (CHILD’S NAME), and repeat what he/she says by adding new words to the conversation. Remember that conversations with you are the greatest treasure for (CHILD’S NAME). 😊
Sentences/Voice tone variations	36+ months	Informative	It is common at this stage of (CHILD’S NAME)'s development to talk to him/her using a higher-pitched, more singsong voice than usual (higher pitched), making each vowel much longer and using short sentences. This way of talking to him/her is good! It helps (CHILD’S NAME) understand the messages you give her better and allows her to identify new words better. ✨
		Activity proposal	Today we propose that at the time of bathing (CHILD’S NAME), you tell him/her and ask questions about what is around using variations in the tone of your voice. 🗣️💧 For example, "Where is the soap; what are we going to dry ourselves with; what do we wrap our feet with?"
		Reflection	How have you felt this week practicing pitch variations in your voice? Variations in the pitch of your voice help (CHILD’S NAME) be more attentive to the sounds of language, convey affection and allow your child to locate where each word begins and ends in a sentence. It generally comes naturally to us to talk to our children this way - don't hold it back, keep doing it, it's very good! 🗣️😊
Music time	6-12 months	Informative	Children recognize their parents' voices from the time they are in the womb. Parents' singing caresses without touching, soothes without speaking and reassures without being so close. When you sing to (CHILD’S NAME), you are helping them develop understanding, use of language and memory. 🎵 If you opt for rhyming songs, the benefits are even greater: you get (CHILD’S NAME) to identify familiar sounds, and that will help him/her when he/she starts reading and spelling. 😊

			We share some links with songs: [LINK TO SONG PLAYLIST]
		Activity proposal	Today we invite you to choose a song that you and (CHILD'S NAME) like. Sing in a tender and loving way, pause to watch (CHILD'S NAME)'s expression and sing again. This can be at any time of the day: when waking up, during playtime, feeding, bath time or before going to sleep. And if your child associates a song to each of these stages, so much the better. The song will allow him/her to anticipate the routine ahead. 🥰
		Reflection	How was this week of singing with your child? Music produces substances in the brain that send feelings of well-being to our bodies. When you and your child feel good, communication with each other and with others improves. Music strengthens the emotional bond that unites you with (CHILD'S NAME). Look for these moments every day! 🎵 🤗
Reading	24-36 months	Informative	When you tell stories or read stories to (CHILD'S NAME), you are teaching them to listen, to concentrate, to understand language, to memorize and to express themselves. All of these skills are essential to his development. Also, sitting close to him/her while you share the story or pictures, strengthens the bond between (CHILD'S NAME) and you. 📖 🌈 We share a page where you can find some stories: [LINK]
		Activity proposal	Today we invite you to share a moment of reading with (CHILD'S NAME). 📖 If you don't have a children's book, you can resort to the website we gave you yesterday, tell him/her a story you remember or make one up. Unfold your creativity, and let the moment flow. Exaggerate gestures, voice tones and body movements to capture (CHILD'S NAME)'s attention, giving him/her space to participate as well. Also try using open-ended questions that arouse his/her curiosity. For example, you can say: "What do you think will happen next?" 💬 Repeat this reading moment every day as a routine. ❤️
		Reflection	How did you do this week in trying to find story time with your child? Each book you read or story you tell is like a brick on which (CHILD'S NAME) will build. The more bricks, the more solid and secure the foundation. Try to make time for these moments every day. What you do for your child today will help him/her forever. 🌈

Note: messages were sent according to the age of the child. In the table we show a selection of messages sent using different age ranges as an example.

Figure A1: Viewing messages in WhatsApp.





Figure A2: Viewing the chatbot for treated families on WhatsApp.



Appendix B: Recruitment

The initial database contained 4436 records of children.

The recruitment of participating families consisted of two stages:

1. First stage (October-November 2021): a first contact call was made to verify that families had a child under three years of age at home and that they were interested in participating in the program. We could contact 35% of the initial database. From these, 86% showed interest in participating in the program. 256 families did not have a child between ages 0-3 and were excluded from the sample.
2. Second stage (December 2021-January 2022): those families who showed interest in participating in the program and who met the entry requirements were contacted again.¹⁹ A baseline survey was applied, and participation consents were recorded. We could survey 79% of contacted households.

In cases where there was more than one child under 3 years of age in the home, the baseline survey was carried out for the youngest child. In the case of twins, two forms were applied per household, one for each child, although we then randomly selected one twin to participate in the evaluation.

Given that the sample size obtained in the second stage was lower than the objective (N=900), a new period of baseline information collection was planned between February and March 2022. On that occasion, MIDES provided a new list of families with a profile similar to families on the PAF waiting list. In particular, the list included families that, due to their characteristics, met the criteria to receive the Uruguay Social Card (TUS), but were not yet receiving it.²⁰

In the new survey period, the surveys were administered by MIDES teleoperators. It was possible to survey 264 households. Adding these surveys, the number of project participants stood at 917 households.

Lastly, in May and July 2022, MIDES provided two additional databases with new families on the waiting list for the PAF. 109 new families were added to the 917 that were already part of the project, bringing the total sample size to 1,026 households.

¹⁹ Children 36 months or younger as of December 6, 2021.

²⁰ Only very poor families are eligible to receive TUS.

Appendix C: Compliance

Table C1: Descriptive statistics of treated families opted out from the program

	Treated families that opted out			Treated families that did not opt out			Difference		
	N	Mean	s.d.	N	Mean	s.d.	N	Coefficient	s.e.
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
<i>Panel A: Child</i>									
Age (months)	57	27413	11374	794	27107	12142	851	0.307	(1.658)
Male	57	0.368	0.487	794	0.525	0.500	851	-0.157**	(0.068)
First son	56	0.607	0.493	791	0.497	0.500	847	0.110	(0.069)
Twin	57	0.000	0.000	794	0.042	0.200	851	-0.042	(0.026)
Attends an early childhood center	57	0.404	0.495	792	0.491	0.500	849	-0.088	(0.069)
Lives with mother	57	0.965	0.186	792	0.981	0.136	849	-0.016	(0.019)
Father or male caregiver in hhld	57	0.596	0.495	794	0.620	0.486	851	-0.023	(0.067)
<i>Panel B: Caregiver</i>									
Woman	57	1000	0.000	794	0.987	0.112	851	0.013	(0.015)
Age (years)	57	26472	9203	792	28519	36351	849	-2.047	(4.828)
Works	56	0.339	0.478	791	0.240	0.427	847	0.099*	(0.060)
Relationship=mother	57	0.947	0.225	794	0.951	0.216	851	-0.004	(0.030)
Completed middle school	57	0.351	0.481	794	0.321	0.467	851	0.030	(0.064)
Risk of depression	57	0.281	0.453	793	0.372	0.484	850	-0.091	(0.066)
Frequency with no internet access = never	45	0.356	0.484	589	0.324	0.469	634	0.031	(0.073)
Frequency with no internet access = once a month	45	0.111	0.318	589	0.160	0.367	634	-0.048	(0.056)
Frequency with no internet access = every 15 days	45	0.222	0.420	589	0.267	0.443	634	-0.044	(0.068)
Frequency with no internet access = every week	45	0.311	0.468	589	0.250	0.433	634	0.062	(0.067)
<i>Panel C: Household</i>									
Number of members	57	4298	1964	793	4738	2222	850	-0.439	(0.302)
Number of children	57	1772	1180	792	2061	1206	849	-0.289*	(0.165)
Overcrowding	57	0.684	0.469	793	0.677	0.468	850	0.007	(0.064)
Government assistance	57	0.965	0.186	791	0.952	0.214	848	0.013	(0.029)
Receives TUS	57	0.579	0.498	791	0.531	0.499	848	0.048	(0.068)
Receives AFAM-PE	57	0.789	0.411	791	0.783	0.413	848	0.007	(0.057)
Experienced negative shock	56	0.839	0.371	785	0.842	0.365	841	-0.003	(0.051)
Number of negative shocks	56	1893	1216	785	2097	1515	841	-0.204	(0.207)
Montevideo	57	0.281	0.453	794	0.343	0.475	851	-0.062	(0.065)

Table C2: Descriptive statistics of treated families that were excluded from the program

	Treated families that were excluded			Treated families that were not excluded			Difference		
	N	Mean	s.d.	N	Mean	s.d.	N	Coefficient	s.e.
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
<i>Panel A: Child</i>									
Age (months)	52	23.240	11.574	799	27.380	12.082	851	-4.140**	(1.725)
Male	52	0.500	0.505	799	0.516	0.500	851	-0.016	(0.072)
First son	51	0.471	0.504	796	0.506	0.500	847	-0.036	(0.072)
Twin	52	0.000	0.000	799	0.041	0.199	851	-0.041	(0.028)
Attends an early childhood center	52	0.365	0.486	797	0.493	0.500	849	-0.128*	(0.071)
Lives with mother	52	0.962	0.194	797	0.981	0.136	849	-0.020	(0.020)
Father or male caregiver in hhld	52	0.462	0.503	799	0.628	0.484	851	-0.167**	(0.069)
<i>Panel B: Caregiver</i>									
Woman	52	0.942	0.235	799	0.991	0.093	851	-0.049***	(0.015)
Age (years)	52	30.680	13.160	797	28.232	36.165	849	2.449	-5.039
	52	0.212	0.412	795	0.249	0.433	847	-0.038	(0.062)

Works	52	0.827	0.382	799	0.959	0.199	851	-0.132***	(0.031)
Relationship=mother	52	0.288	0.457	799	0.325	0.469	851	-0.037	(0.067)
Completed middle school	51	0.373	0.488	799	0.365	0.482	850	0.007	(0.070)
Risk of depression	38	0.263	0.446	596	0.331	0.471	634	-0.067	(0.079)
Frequency with no internet access = never	38	0.211	0.413	596	0.153	0.360	634	0.058	(0.061)
Frequency with no internet access = once a month	38	0.342	0.481	596	0.258	0.438	634	0.084	(0.074)
Frequency with no internet access = every 15 days	38	0.184	0.393	596	0.258	0.438	634	-0.074	(0.073)
Frequency with no internet access = every week	52	4.731	1.931	798	4.707	2.225	850	0.024	(0.316)
<i>Panel C: Household</i>	52	2.173	1.248	797	2.033	1.204	849	0.140	(0.173)
Number of members	52	0.538	0.503	798	0.687	0.464	850	-0.148**	(0.067)
Number of children	52	0.885	0.323	796	0.957	0.202	848	-0.073**	(0.030)
Overcrowding	52	0.404	0.495	796	0.543	0.498	848	-0.139*	(0.071)
Government assistance	52	0.635	0.486	796	0.793	0.406	848	-0.158***	(0.059)
Receives TUS	50	0.860	0.351	791	0.841	0.366	841	0.019	(0.053)
Receives AFAM-PE	50	2.360	1.687	791	2.066	1.484	841	0.294	(0.218)
Experienced negative shock	52	0.269	0.448	799	0.343	0.475	851	-0.074	(0.068)
Number of negative shocks	52	23.240	11.574	799	27.380	12.082	851	-4.140**	(1.725)
Montevideo	52	0.500	0.505	799	0.516	0.500	851	-0.016	(0.072)

Appendix D: Evaluation scales

Table D1: Description of variables included in follow-up questionnaires.

Category	Scale	Follow-up	Batch
	Parental knowledge: 4 questions that survey parents' knowledge of positive parenting. The questions are True/False.	Follow-up 1, Follow-up 2	Batches 1 and 2
	StimQ: StimQ is a scale developed, reviewed and validated by pediatricians and developmental psychologists at New York University. To measure frequency of involvement in didactic activities, we use four StimQ questions that are answered on a scale from 0 to 7 according to the caregiver's frequency with the child in different activities. To measure quality of interaction in didactic activities, we added questions that are answered dichotomously with Yes/No depending on whether the caregiver performs certain practices that are considered to add quality to language interactions such as describing the environment, putting into words what the adult and child are doing, among others.	Follow-up 2	Batches 1 and 2

Depression, Stress, and Wellbeing of caregiver	CESD 10: Center for Epidemiologic Studies scale used to measure depression. It consists of 10 questions and uses a 4-option Likert scale, where 1 indicates that the respondent rarely or never experienced the symptoms described, and 4 indicates that the symptom was experienced most days or every day. A depression risk indicator is constructed that takes the value of 1 if the score is equal to or greater than 10.	Follow-up 1, Follow-up 2	Batches 1 and 2
	Parental stress: We applied the dysfunctional parent-child interaction subscale of the Parental Stress Index (PSI). This is a self-reported questionnaire that allows us to identify different sources of stress that emerge during parenting. The subscale we use approximates the degree of satisfaction that the caregiver has with his or her child and with the interactions they have. The respondent must answer how much he/she agrees with certain statements referring to common parenting situations on a five-point Likert scale. Additionally, the PSI scale incorporates a question that seeks to survey how the caregiver perceives him/herself (e.g. "Not a very good caregiver", "A person who has some problems being a caregiver", "A normal caregiver", "A better than average caregiver", "A very good caregiver").	Follow-up 1, Follow-up 2	Batch 1 (second follow-up), Batch 2 (first and second follow-up)
	WHO-5: WHO-5 of the World Health Organization (WHO), a self-reported questionnaire that measures current mental well-being. The WHO-5 has demonstrated adequate validity as a screening instrument for depressive symptoms. It consists of five questions regarding vitality, mood and general interest answered on a six-point scale. The time reference of the instrument is the last two weeks. Based on the score obtained, a risk threshold of depression is established.	Follow-up 1, Follow-up 2	Batch 1 (second follow-up), Batch 2 (first and second follow-up)
	McConkey Well-Being Scale (2020): An instrument based on the concept of subjective well-being that seeks to measure how satisfied the person is in various dimensions of his or her life (health, household and care tasks, general quality of life, etc.) on a 10-point scale.	Follow-up 1, Follow-up 2	Batch 1 (second follow-up), Batch 2 (first and second follow-up)

	We constructed a well-being index by averaging the score on two items from the reduced version of the Burden Scale for Family Caregivers (BSFC) and four items from the Well-Being During Pregnancy (WiP) questionnaire developed by Alderdice et al. (2017) and Kelly et al. (2022) that we adapted to the parenting stage in life. Specifically, we employed the following statements from the BSFC: "Your satisfaction with your life has been negatively affected due to caring for your child" and "Caring for your child has taken quite a bit of your energy." The person should indicate how much he or she agrees or disagrees with the statement on a four-point scale where 1 is strongly disagree and 4 is strongly agree. Lower values indicate less caregiving overload. From the WiP scale we used four items that seek to capture the frequency with which the caregiver feels connected with his/her child, enjoys his/her role as caregiver, has social contact with other people and feels confident in how to care for his/her child.	Follow-up 1, Follow-up 2	Batch 1 (second follow-up), Batch 2 (first and second follow-up)
--	--	---------------------------------	---

Appendix E: Descriptive statistics of the outcome variables

Table E1: Descriptive statistics of the outcome variables for batch 1

	Control			Treatment			Difference		
	N	Mean	s.d.	N	Mean	s.d.	N	Coefficient	s.e.
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
<i>Panel A: Child</i>									
Age (months)	352	25.328	13.202	590	27.206	12.338	942	0.151	(0.693)
Male	352	0.509	0.501	590	0.504	0.501	942	-0.007	(00.034)
First son	352	0.428	0.495	590	0.474	0.500	940	0.055	(00.034)
Twin	352	0.341	0.182	590	0.458	0.210	942	0.007	(00.013)
Attends an early childhood center	352	0.506	0.501	590	0.519	0.501	942	0.021	(00.034)
Lives with mother	352	0.972	0.167	590	0.980	0.142	942	0.006	(0.059)
Father or male caregiver in hhld	352	0.614	0.488	590	0.631	0.484	941	0.007	(0.010)
<i>Panel B: Caregiver</i>							942	0.013	(0.033)
Woman	352	0.998	0.0534	590	0.985	0.123	942	-0.014*	(0.007)
Age (years)	352	28.098	7.722	590	29.316	41.869	939	1.380	(2.278)
Works	352	0.257	0.437	590	0.259	0.438	939	-0.004	(0.030)
Relationship=mother	352	0.944	0.232	590	0.946	0.227	942	0.002	(0.015)
Completed middle school	352	0.296	0.457	590	0.333	0.472	942	0.040	(0.032)
Completed high school	352	0.080	0.271	590	0.051	0.220	942	-0.029*	(0.016)
Risk of depression	352	0.407	0.492	590	0.360	0.480	941	-0.048	(0.033)
Frequency with no internet access = every week	352	0.310	0.401	590	0.320	0.402	704	0.013	(0.036)
Frequency with no internet access = once a month	352	0.168	0.336	590	0.147	0.311	704	-0.019	(0.029)

Frequency with no internet access = every 15 days	352	0.278	0.391	590	0.278	0.387	704	-0.006	(0.035)
Frequency with no internet access = every week	352	0.245	0.371	590	0.256	0.375	704	0.011	(0.034)
<i>Panel C: Household</i>							942	0.134	(0.146)
Number of members	352	4.787	1.962	590	4.829	2.321	937	-0.098	(0.084)
Number of children	352	2.171	1.374	590	2.048	1.159	937	-0.022	(0.013)
Overcrowding	352	0.676	0.469	590	0.672	0.471	937	-0.041	(0.034)
Government assistance	352	0.974	0.159	590	0.951	0.217	937	-0.002	(0.027)
Receives TUS	352	0.593	0.490	590	0.541	0.498	942	-0.051*	(0.030)
Receives AFAM-PE	352	0.799	0.400	590	0.778	0.416	927	0.003	(0.025)
Experienced negative shock	352	0.850	0.355	590	0.845	0.360	927	0.043	(0.103)
Number of negative shocks	352	2.120	1.483	590	2.140	1.514	896	-0.006	(0.007)
Montevideo	352	0.350	0.478	590	0.339	0.474	941	-0.001	(0.032)

Table E2: Descriptive statistics of the outcome variables for batch 2

	Control			Treatment			Difference		
	N	Mean	s.d.	N	Mean	s.d.	N	Coefficient	s.e.
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
<i>Panel A: Child</i>									
Age (months)	269	25.359	12.921	484	27.784	12.279	753	1.083	(0.774)
Male	269	0.506	0.501	484	0.515	0.501	753	0.010	(0.038)
First son	269	0.439	0.498	484	0.485	0.500	752	0.051	(0.038)
Twin	269	0.034	0.181	484	0.046	0.209	753	0.009	(0.015)
Attends an early childhood center	269	0.517	0.501	484	0.515	0.500	751	0.006	(0.038)

Lives with mother	269	0.978	0.148	484	0.978	0.150	753	0.029	(0.068)
Father or male caregiver in hhld	269	0.621	0.487	484	0.643	0.480	751	0.000	(0.011)
<i>Panel B: Caregiver</i>							753	0.019	(0.037)
Woman	269	1	0	484	0.996	0.065	753	-0.005	(0.004)
Age (years)	269	27.855	7.541	484	27.516	8.164	751	-0.547	(0.604)
Works	269	0.243	0.429	484	0.261	0.440	751	0.017	(0.033)
Relationship=mother	269	0.960	0.199	484	0.961	0.195	753	0.003	(0.015)
Completed middle school	269	0.302	0.460	484	0.362	0.481	753	0.061*	(0.036)
Completed high school	269	0.082	0.275	484	0.052	0.222	753	-0.031*	(0.018)
Risk of depression	269	0.387	0.488	484	0.375	0.485	752	-0.012	(0.037)
Frequency with no internet access = every week	269	0.338	0.406	484	0.321	0.396	545	-0.027	(0.042)
Frequency with no internet access = once a month	269	0.147	0.308	484	0.140	0.298	545	-0.003	(0.032)
Frequency with no internet access = every 15 days	269	0.280	0.385	484	0.282	0.384	545	-0.002	(0.041)
Frequency with no internet access = every week	269	0.237	0.358	484	0.259	0.371	545	0.032	(0.039)
<i>Panel C: Household</i>							752	0.215	(0.162)
Number of members	269	4.614	1.802	484	4.780	2.292	750	-0.024	(0.093)
Number of children	269	2.090	1.290	484	2.050	1.174	748	-0.014	(0.014)
Overcrowding	269	0.654	0.477	484	0.684	0.466	748	-0.044	(0.038)
Government assistance	269	0.978	0.148	484	0.963	0.190	748	0.004	(0.031)
Receives TUS	269	0.581	0.493	484	0.526	0.499	752	-0.048	(0.034)
Receives AFAM-PE	269	0.787	0.410	484	0.776	0.417	742	-0.005	(0.027)
Experienced negative shock	269	0.857	0.349	484	0.849	0.356	742	-0.077	(0.116)
Number of negative shocks	269	2.176	1.580	484	2.077	1.450	714	-0.003	(0.008)
Montevideo	269	0.370	0.483	484	0.352	0.478	751	0.034	(0.036)

Table E3: Descriptive statistics of the outcome variables for batch 3

	Control			Treatment			Difference		
	N	Mean	s.d.	N	Mean	s.d.	N	Coefficient	s.e.
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
<i>Panel A: Child</i>									
Age (months)	126	25.385	14.398	186	27.855	12.676	1.835	(1.243)	312
Male	126	0.397	0.492	186	0.587	0.494	0.188***	(0.057)	312
First son	126	0.437	0.498	186	0.441	0.498	0.009	(0.057)	312
Twin	126	0.040	0.196	186	0.027	0.163	-0.013	(0.020)	312
Attends an early childhood center	126	0.532	0.501	185	0.584	0.495	0.057	(0.057)	311
Lives with mother	125	0.992	0.090	186	0.974	0.163	-0.018	(0.016)	311
Father or male caregiver in hhld	126	0.659	0.477	186	0.630	0.485	-0.031	(0.056)	312
<i>Panel B: Caregiver</i>									
Woman	126	1	0	186	0.995	0.074	-0.006	(0.007)	312
Age (years)	126	28.286	7.203	185	29.160	8.487	0.691	(0.911)	311
Works	125	0.224	0.419	185	0.255	0.437	0.035	(0.050)	310
Relationship=mother	126	0.969	0.177	186	0.952	0.216	-0.014	(0.023)	312
Completed middle school	126	0.326	0.471	186	0.361	0.482	0.037	(0.055)	312
Completed high school	126	0.088	0.284	186	0.060	0.237	-0.031	(0.030)	312
Risk of depression	126	0.381	0.488	185	0.384	0.488	0.010	(0.056)	311
Frequency with no internet access = every week	98	0.327	0.472	141	0.235	0.425	-0.096	(0.059)	239
Frequency with no internet access = once a month	98	0.103	0.305	141	0.192	0.395	0.093*	(0.047)	239
Frequency with no internet access = every 15 days	98	0.296	0.459	141	0.320	0.468	0.019	(0.061)	239

Frequency with no internet access = every week	98	0.276	0.450	141	0.256	0.438	-0.016	(0.058)	239
<i>Panel C: Household</i>									
Number of members	126	4.747	1.753	185	4.455	1.859	-0.280	(0.210)	311
Number of children	126	2.254	1.486	186	1.968	1.120	-0.276*	(0.148)	312
Overcrowding	126	0.683	0.468	185	0.687	0.466	0.011	(0.054)	311
Government assistance	126	0.993	0.089	186	0.963	0.191	-0.029	(0.018)	312
Receives TUS	126	0.651	0.479	186	0.565	0.498	-0.079	(0.057)	312
Receives AFAM-PE	126	0.802	0.401	186	0.807	0.397	0.015	(0.045)	312
Experienced negative shock	123	0.846	0.363	184	0.870	0.338	0.029	(0.041)	307
Number of negative shocks	123	2.383	1.791	184	2.191	1.480	-0.147	(0.184)	307
Montevideo	126	0.366	0.484	186	0.269	0.445	-0.106**	(0.051)	312

Table E4 shows descriptive statistics of the outcome variables for the control families on the follow-up questionnaires. 61% and 64% of those surveyed receive the Uruguay Social Card (TUS) in the first and second follow-up respectively. 55% receive the Parenting Bonus²¹ in the first follow-up and 49% receive it in the second. 19% and 21% of households obtained information about work programs in the four months prior to the first and second follow-up respectively. 6% of households accessed a program to improve their home in the months prior to the first follow-up and 5% in the months prior to the second.

Regarding food support in the 4 months prior to the first survey, 8% of households received food baskets in the first follow-up and 10% in the second, and 17% and 19% started paperwork to receive food support in the first and second follow-up respectively.

32% of the caregivers reported working during the first follow-up and 38% during the second. 65% of the children attend an early childhood center and 78% do so at the end of the intervention. Children who receive free early childhood vouchers (BIS, for its acronym in Spanish: *Beca de inclusión socio-educativa*) are 3% of the sample in the first follow-up and 4% in the second.

Regarding parental involvement in cognitive stimulation activities, on the Stim-Q scale, families average a value of 4.9 on a scale that goes from 0 to 7. Regarding the quality of language, families score 1.1 with a deviation of 0.18 on a scale that goes from 1 to 1.875. On the parental knowledge scale, the caregivers answered on average 2.5 and 2.6 of 4 questions correctly in the first and second follow-up respectively. On the parental knowledge scale in language topics, parents respond 2.2 out of 3 questions correctly on average.

Regarding the emotional well-being of the caregiver, the risk of depression is 50% in the first follow-up and 54% in the second follow-up. This is consistent with the fact that families that opt out of the program are less likely to present risk of depression at baseline (see Table C1 in Appendix C). Parental stress has a score of 21 points. On the WHO-5 scale, whose score ranges from 0 to 25, the average was 16 points in both follow-ups. On the Mc Conkey's (2000) subjective well-being scale, families scored an average of 62 in the first and second follow-up, out of a total of 80. In the wellbeing index we constructed, that can take values between 2.67 and 6.67, control families have a score of 5 on average in the first follow-up and in the second follow-up.

Regarding anthropometry, height for age and wage for age is below the standardized mean (negative values) in both follow-ups.

²¹ The Parenting Bonus is implemented through a monthly monetary transfer that is paid through the Uruguay Social Card (TUS) it is aimed at households in situations of extreme socioeconomic vulnerability in which pregnant women and/or children from 0 to 3 reside. years.

Table E4: Descriptive statistics of the outcome variables for the control families in the follow-up questionnaires

Variable	First follow-up					Second follow-up				
	N	Mean	s.d.	Min	Max	N	Mean	s.d.	Min	Max
Panel A: Government transfers and programs										
Government transfers	323	-0.233	1.255	-2.011	2.005	256	-0.258	1.244	-2.011	2.302
Uruguay Social Card (TUS)	351	0.613	0.488	0	1	268	0.642	0.480	0	1
Parenting Bonus	332	0.545	0.499	0	1	261	0.490	0.501	0	1
Labor Programs	350	0.194	0.396	0	1	266	0.214	0.411	0	1
Housing Improvement	350	0.060	0.238	0	1	268	0.049	0.215	0	1
Food baskets	348	0.083	0.277	0	1	264	0.102	0.304	0	1
Started paperwork for food program	340	0.168	0.374	0	1	266	0.188	0.391	0	1
Caregiver works	351	0.319	0.467	0	1	268	0.381	0.486	0	1
Child attends early childhood center	352	0.653	0.477	0	1	268	0.776	0.418	0	1
Child receives BIS voucher to attend center	350	0.029	0.167	0	1	267	0.041	0.199	0	1
Panel B: Parental investment and knowledge										
Stim-Q						265	4.870	1.346	0	7
Quality language stimulation						265	1.125	0.178	1	1.875
Parental knowledge - childrearing	350	2.517	1.056	0	4	262	2.649	1.005	0	4
Parental knowledge - language						260	2.238	0.723	0	3

Panel C: Parental Wellbeing and Stress

Wellbeing PCA	190	-0.001	1.514	-3.872	2.832	246	-0.142	1.501	-3.815	2.976
Risk of depression	352	0.497	0.501	0	1	263	0.540	0.499	0	1
WHO-5	196	16.082	5.549	1	25	258	15.508	5.452	0	25
McConkey Well-being Scale (2000)	192	62.078	12.426	29	80	252	61.687	11.726	27	80
Wellbeing index	195	5.140	0.788	2.667	6.667	255	5.114	0.842	2.667	6.667
Parental stress	197	21.355	7.016	12	45	252	20.647	7.444	12	45
Panel D: Child's anthropometry										
Height for age	128	-1.209	1.122	-3.404	2.210	139	-1.118	1.459	-4.973	4.136
Weight for age	138	-0.321	1.043	-4.156	1.960	149	-0.259	1.154	-4.411	2.962
Panel E: Language outcomes										
Mother's average word per minute	175	251.798	60.946	123.045	485.887	125	246.187	53.352	168.045	464.903
Mother's vocalizations per hour	176	891.107	234.985	0	1622.352	126	900.203	236.203	0	1651.477
Mother's vocalizations duration	176	1.73e+06	6.25e+05	0	3.34e+06	126	1.73e+06	5.52e+05	0	2.75e+06
Child's vocalizations per hour	176	672.841	306.766	0	1333.580	126	752.969	258.317	83.772	1541.379
Child's vocalizations duration	176	8.11e+05	4.99e+05	0	2.99e+06	126	9.96e+05	5.34e+05	44255.330	3.26e+06
Pitch range	175	23.215	3.189	12.393	31.649	125	23.074	2.777	9.828	30.019
Mother's average pitch	175	24.082	2.370	16.882	30.053	125	24.223	2.142	17.456	30.676
Child-adult conversational transitions	176	797.641	447.640	0	2089.671	126	924.510	437.489	65.291	2862.560

Appendix F: Full set of results

Table F1 shows results for the full set of outcomes (used to construct summary indices). Regarding the effect of the program on access to monetary transfers, we see that in the first follow-up the treated have greater access to the TUS and the Parenting Bonus. The effects are of a magnitude of 0.26 and 0.2 standard deviations respectively. In the second follow-up, the effects are of a magnitude of 0.36 standard deviations for TUS and 0.18 standard deviations for Parenting Bonus. There are no significant differences between treated and controls in access to the Housing Improvement Program, but the power to detect significant effects in this outcome is very low (see Table G1 in Appendix G).

We find that, on average, those treated have accessed more food baskets than controls in the 30 days prior to the first follow-up survey (Panel C). The magnitude of the effect is of 0.31 standard deviations. These baskets were awarded punctually and at the discretion of the teleoperator. In the second follow-up, after the implementation of the module exclusively oriented to language development, we did not find significant differences between treated and controls in access to food support.

With respect to attendance at educational centers, we see that in the first follow-up, children from treated families attend early childhood centers to a greater extent than controls, but in the second follow-up this effect disappears. The receipt of BIS scholarships is also higher among treated children.

In terms of parental wellbeing, we find that there is a positive effect on the WHO-5 score and index of wellbeing that is significant in the second follow-up.

Table F1: Effects of the intervention on access to government programs, parental involvement, caregiver well-being and satisfaction with the program.

	First follow-up (1)	Second follow-up (2)
Uruguay Social Card	0.257*** (0.066) 939	0.364*** (0.072) 746
Parenting Bonus	0.199*** (0.066) 897	0.178** (0.076) 725
Home improvement programs	-0.032 (0.067) 938	-0.068 (0.073) 749
Food baskets	0.305*** (0.079) 927	-0.040 (0.076) 739
Paperwork for food basket	0.101 (0.071) 900	-0.137* (0.074) 740
Caregiver works	0.051 (0.068) 939	-0.028 (0.076) 751
Child attends early childhood center	0.131** (0.066) 941	-0.052 (0.077) 752
BIS scholarship	0.144* (0.076) 935	0.046 (0.080) 745
Risk of depression	0.004 (0.068) 937	-0.097 (0.077) 737
WHO 5 wellbeing	0.064 (0.095) 482	0.134* (0.076) 726
Wellbeing score	0.075 (0.092) 472	0.112 (0.080) 707
Index of wellbeing	-0.128 (0.099) 476	0.138* (0.078) 724

Notes: * $p < .1$, ** $p < .05$, *** $p < .01$. Rows depict different outcomes. Families of outcomes are identified by a heading in italics. Columns (1) and (5) report the ITT coefficient (outcome difference between ITT=1 and ITT=0) in the first and second follow-up respectively. Columns (2)-(3) and (6)-(7) report the ITT coefficient of specifications that include controls. We adjust for multiple hypothesis testing the p-values of the coefficients in columns (1) and (5) in columns (4) and (8) respectively. Families of outcomes are marked

with a heading in bold and italics. In families with only one outcome, we do not adjust p-values for multiple hypothesis testing and we mark those cases with n/a. Number of observations in straight parentheses.

Appendix G: Power calculations

Table G1: Power calculations to detect an effect of 0.2 standard deviations in continuous variables and of 20 percentage points in dichotomous variables

	First follow-up (1)	Second follow-up (2)
<i>Monetary transfers</i>		
Government transfers	0.838	0.753
Uruguay Social Card	0.973	0.960
Parenting Bonus (up to 3 years of age)	0.898	0.722
Food baskets (in the last 30 days)	0.129	0.127
Home improvement program	0.104	0.081
Child receives BIS voucher to attend center	0.071	0.075
Labor programs (duration 6 months)	0.282	0.254
<i>Parental inputs</i>		
Parental knowledge – Childrearing	0.842	0.724
Parental knowledge – Language		0.758
StimQ - Frequency		0.754
StimQ - Quality		0.784
<i>Parental stress and wellbeing</i>		
Wellbeing PCA	0.577	0.720
Depression (CESD $\geq$ 10)	0.844	0.819
WHO-5	0.581	0.742
Caregiver well-being: Mc Conkey (2000)	0.586	0.718
Wellbeing index	0.533	0.731
Parenting Stress Index	0.574	0.734
<i>Child's anthropometry</i>		
Height for age	0.369	0.471
Weight for age	0.352	0.470

Note: The power calculations consider an effect size of 0.20 standard deviations in the case of continuous outcomes and of 20 percentage points in the case of dichotomous outcomes.

Bibliography

Aizer, A. (2003). Low take-up in Medicaid: Does outreach matter and for whom? *American Economic Review*, 93(2), 238-241.

Ajzenman, N., Luna, L. B., Hernández-Agramonte, J. M., Boo, F. L., Alfaro, M. P., Vásquez-Echeverría, A., & Diaz, M. M. (2022). A behavioral intervention to increase preschool attendance in Uruguay. *Journal of Development Economics*, 159, 102984.

Arteaga, I., Trias, J., 2023. Can technology narrow the early childhood stimulation gap in rural Guatemala? Results from an experimental approach. *Unpublished manuscript*. Washington, DC: The World Bank.

Arteaga, I., de Barros, A., & Ganimian, A. J. (2024). The Challenges of Scaling up Effective Child-Rearing Practices Using Technology in Developing Settings: Experimental Evidence From India.

Attanasio, O., Baker-Henningham, H., Bernal, R., Meghir, C., Pineda, D., & Rubio-Codina, M. (2022). Early stimulation and nutrition: the impacts of a scalable intervention. *Journal of the European Economic Association*, 20(4), 1395-1432.

Alderdice, F., McNeill, J., Gargan, P., & Perra, O. (2017). Preliminary evaluation of the Well-being in Pregnancy (WiP) questionnaire. *Journal of Psychosomatic Obstetrics & Gynecology*, 38(2), 133-142.

Almond, D., Currie, J., & Duque, V. (2018). Childhood circumstances and adult outcomes: Act II. *Journal of Economic Literature*, 56(4), 1360-1446.

Balsa, A., Boo, F. L., Bloomfield, J., Cristia, A., Cid, A., de La Paz Ferro, M., ... & del Lujan Gonzalez, M. (2021). Effect of Crianza Positiva e-messaging program on adult-child language interactions. *Behavioural Public Policy*, 1-37.

Balsa, A., Bloomfield, J., & Cid, A. (2022). The Replication of a Parenting Behavioral Change Communication Intervention during the COVID-19 Pandemic: Too Much or Too Little Information? Working paper, Universidad de Montevideo. <https://n9.cl/elk7u>

Balsa, A., Bloomfield, J., Cid, A., Lorenzo, M., & Patrone, P. (2023a). Household's stressors and well-being during the global COVID-19 pandemic. *Journal of Family Issues*, 0192513X231194289.

Balsa, A. I., Gómez Muzzio, E., González, M. L., Bloomfield, J., Cid, A., & Valdés, R. (2023b). Crianza positiva: combining group workshops and e-messages to strengthen parenting competences. In *Child & Youth Care Forum* (pp. 1-22). New York: Springer US.

Berlinski, S., & Schady, N. (2015). *The early years: child well-being and the role of public policy* (p. 288). Springer Nature.

Bhargava, S., & Manoli, D. (2015). Psychological frictions and the incomplete take-up of social benefits: Evidence from an IRS field experiment. *American Economic Review*, 105(11), 3489-3529.

Bloomfield, J., Balsa, A., & Cid, A. (2022). Using behavioral insights in early childhood interventions: the effects of Crianza Positiva e-messaging program on parental investment. *Review of Economics of the Household*, 1-36.

Bradley, R. H., Corwyn, R. F., McAdoo, H. P., & García Coll, C. (2001). The home environments of children in the United States part I: Variations by age, ethnicity, and poverty status. *Child development*, 72(6), 1844-1867.

Britto, P. R., Lye, S. J., Proulx, K., Yousafzai, A. K., Matthews, S. G., Vaivada, T., ... & Bhutta, Z. A. (2017). Nurturing care: promoting early childhood development. *The Lancet*, 389(10064), 91-102.

Cabrera, N. J., Shannon, J. D., Vogel, C., Tamis-LeMonda, C., Ryan, R. M., Brooks-Gunn, J., Raikes, H., & Cohen, R. (2004). Low-income fathers' involvement in their toddlers' lives: biological fathers from the early head start research and evaluation study. *Fathering: A Journal of Theory, Research & Practice About Men as Fathers*, 2(1), 5–30.

Currie, J. (2004). The take up of social benefits. *National Bureau of Economics Research*.

Doepke, M., Sorrenti, G., & Zilibotti, F. (2019). The economics of parenting. *Annual Review of Economics*, 11, 55-84.

Doss, Christopher, Erin M. Fahle, Susanna Loeb, and Benjamin N. York. 2017. “Supporting Parenting through Differentiated and Personalized Text-Messaging: Testing Effects on Learning during Kindergarten.” CEPA Working Paper 16-18. Stanford Center for Education Policy Analysis.

Evans, William D., Jasmine L. Wallace, and Jeremy Snider. 2012. Pilot evaluation of the text4baby mobile health program. *BMC Public Health* 12(1): 1031.

Ferjan Ramírez, N., S. R. Lytle, M. Fish, and P. K. Kuhl (2019). Parent coaching at 6 and 10 months improves language outcomes at 14 months: A randomized controlled trial. *Developmental Science* 22 (3), e12762.

Ferjan Ramírez, N., S. R. Lytle, and P. K. Kuhl (2020). Parent coaching increases conversational turns and advances infant language development. *Proceedings of the National Academy of Sciences* 117 (7), 3484–3491.

Gallego, F., Oreopoulos, P., & Spencer, N. (2023). *The Importance of a Helping Hand in Education and in Life* (No. w31706). National Bureau of Economic Research.

Gertler, P., Heckman, J., Pinto, R., Zanolini, A., Vermeersch, C., Walker, S., ... & Grantham-McGregor, S. (2014). Labor market returns to an early childhood stimulation intervention in Jamaica. *Science*, 344(6187), 998-1001.

Grantham-McGregor, S., Cheung, Y. B., Cueto, S., Glewwe, P., Richter, L., & Strupp, B. (2007). Developmental potential in the first 5 years for children in developing countries. *The lancet*, 369(9555), 60-70.

Guryan, J., Hurst, E., & Kearney, M. (2008). Parental education and parental time with children. *Journal of Economic perspectives*, 22(3), 23-46.

Heckman, J. J. (2006). Skill formation and the economics of investing in disadvantaged children. *Science*, 312(5782), 1900-1902.

Hurwitz, Lisa B., Alexis R. Lauricella, Ann Hanson, Anthony Raden, and Ellen Wartella. 2015. “Supporting Head Start parents: impact of a text message intervention on parent–child activity engagement.” *Early Child Development and Care* 185(9):1373–1389.

- Kelly, L., Kurinczuk, J. J., Fitzpatrick, R., & Alderdice, F. (2022). Refinement of the Well-being in Pregnancy (WiP) questionnaire: cognitive interviews with women and healthcare professionals and a validation survey. *BMC pregnancy and childbirth*, 22(1), 325.
- Law, J., Rush, R., Schoon, I., & Parsons, S. (2009). Modeling developmental language difficulties from school entry into adulthood: Literacy, mental health, and employment outcomes.
- Leech, K., R. Wei, J. R. Harring, and M. L. Rowe (2018). A brief parent-focused intervention to improve preschoolers' conversational skills and school readiness. *Developmental Psychology* 54 (1), 15.
- Leech, K. A. et al. (2021). An intervention to increase conversational turns between parents and young children. *Journal of Child Language* 48 (2), 399–412.
- Leung, C. Y., M. W. Hernandez, and D. L. Suskind (2020). Enriching home language environment among families from low-ses backgrounds: A randomized controlled trial of a home visiting curriculum. *Early Childhood Research Quarterly* 50, 24–35.
- Mayer, Susan E., Ariel Kalil, Philip Oreopoulos, and Sebastián Gallegos. 2018. Using behavioral insights to increase parental engagement: The parents and children together intervention. *Journal of Human Resources* 0617–8835R.
- Mcconkey, Roy. (2020). A Brief Measure of Parental Wellbeing for Use in Evaluations of Family-Centred Interventions for Children with Developmental Disabilities. *Children*. 7. 10.3390/children7090120.
- McGillion, M., J. M. Pine, J. S. Herbert, and D. Matthews (2017). A randomised controlled trial to test the effect of promoting caregiver contingent talk on language development in infants from diverse socioeconomic status backgrounds. *Journal of Child Psychology and Psychiatry* 58 (10), 1122–1131
- Meuwissen, Alyssa, Alison Giovanelli, Madelyn Labella, and Amy Susman-Stillman. 2017. "Text2Learn: An Early Literacy Texting Intervention by Community Organizations."
- Oreopoulos, P., Petronijevic, U., Logel, C., & Beattie, G. (2020). Improving non-academic student outcomes using online and text-message coaching. *Journal of Economic Behavior & Organization*, 171, 342-360.
- Peurey, L., Havard, W., Virlet, G., Cao, X., Balsa, A., Cid, A., Ottavianelli, M., Horta, J., Cristia, A. (2023). An Automated Pipeline for Providing Personalized Feedback Based on Audio Samples.
- Rafla, J., Schwartz, K., Yoshikawa, H., Hilgendorf, D., Ramachandran, A., Khanji, M., Abu Seriah, R., Al Abed, M., Fityan, R., Sloane, P. and Al Aqra, A., Sharawi, T., Molano, A., Foulds, K., Behrman, J.R., Wuermli, A., 2024. Cluster randomized controlled trial of a phone-based caregiver program for Syrian and Jordanian families with young children. New York, NY: Global TIES for Children, New York University (NYU).
- Rowe, M. L. (2012). A longitudinal investigation of the role of quantity and quality of child-directed speech in vocabulary development. *Child development*, 83(5), 1762-1774.
- Smith, J.A., Chang, S.M., Brentani, A., Fink, G., Lopez-Boo, F., Torino, B.M., Codina, M.R., Walker, S.P., 2023. A remote parenting program and parent and staff perspectives: A randomized trial. *Pediatrics* 151, e2023060221F.

Suskind, D. L., K. R. Leffel, E. Graf, M. W. Hernandez, E. A. Gunderson, S. G. Sapolich, E. Suskind, L. Leininger, S. Goldin-Meadow, and S. C. Levine (2016). A parent-directed language intervention for children of low socioeconomic status: A randomized controlled pilot study. *Journal of child language* 43 (2), 366–406.

Walker, S. P., Wachs, T. D., Grantham-McGregor, S., Black, M. M., Nelson, C. A., Huffman, S. L., ... & Richter, L. (2011). Inequality in early childhood: risk and protective factors for early child development. *The Lancet*, 378(9799), 1325-1338.

York, Benjamin N., Susanna Loeb, and Christopher Doss. 2019. One Step at a Time The Effects of an Early Literacy Text-Messaging Program for Parents of Preschoolers. *Journal of Human Resources* 54(3):537–566.

Zimmerman, F. J., Gilkerson, J., Richards, J. A., Christakis, D. A., Xu, D., Gray, S., & Yipanel, U. (2009). Teaching by listening: The importance of adult-child conversations to language development. *Pediatrics*, 124(1), 342-349.